

BANACH LATTICES WITH UPPER p -ESTIMATES: FREE AND INJECTIVE OBJECTS

E. GARCÍA-SÁNCHEZ, D. H. LEUNG, M. A. TAYLOR, AND P. TRADACETE

ABSTRACT. We study the free Banach lattice $\text{FBL}^{(p,\infty)}[E]$ with upper p -estimates generated by a Banach space E . Using a classical result of Pisier on factorization through $L^{p,\infty}(\mu)$ together with a finite dimensional reduction, it is shown that the spaces $\ell^{p,\infty}(n)$ witness the universal property of $\text{FBL}^{(p,\infty)}[E]$ isomorphically. As a consequence, we obtain a functional representation for $\text{FBL}^{(p,\infty)}[E]$, answering a question from [28]. More generally, our proof allows us to identify the norm of any free Banach lattice over E associated with a rearrangement invariant function space.

After obtaining the above functional representation, we take the first steps towards analyzing the fine structure of $\text{FBL}^{(p,\infty)}[E]$. Notably, we prove that the norm for $\text{FBL}^{(p,\infty)}[E]$ cannot be isometrically witnessed by $L^{p,\infty}(\mu)$ and settle the question of characterizing when an embedding between Banach spaces extends to a lattice embedding between the corresponding free Banach lattices with upper p -estimates. To prove this latter result, we introduce a novel push-out argument, which when combined with the injectivity of ℓ^p allows us to give an alternative proof of the subspace problem for free p -convex Banach lattices. On the other hand, we prove that $\ell^{p,\infty}$ is not injective in the class of Banach lattices with upper p -estimates, elucidating one of many difficulties arising in the study of $\text{FBL}^{(p,\infty)}[E]$.

CONTENTS

1. Introduction	2
1.1. Outline of the paper	4
2. The free Banach lattice associated to a class of Banach lattices	5
2.1. Review of the free vector lattice	5
2.2. Universality of the free Banach lattice associated to a class of Banach lattices	7
2.3. Rearrangement invariant spaces and the norm for $\text{FBL}^{\mathcal{C}}[E]$	8
2.4. Injective homomorphisms need not extend to the norm completion	11
3. Applications to free p -convex and free (p, ∞) -convex Banach lattices	12
3.1. Solution to Problem 2.8 for the classes \mathcal{C}_p and $\mathcal{C}_{p,\infty}$.	16
4. Distinguishing $\rho_{\mathcal{X}_{p,\infty}}$ and $\rho_{\mathcal{C}_{p,\infty}}$	17
4.1. Equivalent renormings of $L^{p,\infty}(\mu)$	21
5. Subspace problem	24
5.1. Push-outs in \mathcal{C}	26
5.2. Recovering the POE- p via injectivity	30
5.3. $\ell^{p,\infty}$ is not injective for Banach lattices with upper p -estimates	32
Acknowledgements	35

Date: March 1, 2024.

2020 Mathematics Subject Classification. 46B42 (primary); 06B25, 47B60 (secondary).

Key words and phrases. Free Banach lattice; upper p -estimate; weak- L^p .

1. INTRODUCTION

Given a Banach space E , the *free Banach lattice generated by E* is a Banach lattice $\text{FBL}[E]$ together with a linear isometric embedding $\phi_E : E \rightarrow \text{FBL}[E]$ such that for every Banach lattice X and every bounded linear operator $T : E \rightarrow X$ there is a unique lattice homomorphism $\widehat{T} : \text{FBL}[E] \rightarrow X$ such that $\widehat{T} \circ \phi_E = T$ and $\|\widehat{T}\| = \|T\|$. In other words, $\text{FBL}[E]$ is the (unique) Banach lattice which converts linear operators defined on E into lattice homomorphisms, in the sense of the following diagram:

$$\begin{array}{ccc} & \text{FBL}[E] & \\ & \uparrow \phi_E & \text{\scriptsize } \exists! \widehat{T} \\ & E & \xrightarrow{T} X \end{array}$$

Over the past ten years, many novel techniques have been developed to understand the structural properties of $\text{FBL}[E]$ as well as the rigidity of the correspondence $E \mapsto \text{FBL}[E]$. This has led to a host of new developments in the theory of Banach spaces and Banach lattices, as well as to the resolution of several outstanding open problems. To indicate just a few examples, free Banach lattices are used in [7] to solve a question of J. Diestel on WCG spaces, in [28] to solve a question from [40] regarding basic sequences satisfying maximal inequalities, and in [14] to produce the first example of a lattice homomorphism which does not attain its norm. Free Banach lattices also play a fundamental role in the theory of projective Banach lattices [4, 5, 15] and are used in [8] to construct Banach lattices of universal disposition, separably injective Banach lattices and push-outs. On the other hand, free Banach lattices are known to have a remarkably rigid structure in their own right. Notably, it is proven in [6] that any disjoint collection in $\text{FBL}[E]$ must be countable and in [28] that $\text{FBL}[E]$ determines E isometrically if E^* is smooth. This latter result has initiated a program to find geometric properties of a Banach space E which can be equivalently characterized via lattice properties of $\text{FBL}[E]$ – see [28] for several such properties.

Although defined in an abstract manner, free Banach lattices admit a very concrete functional representation. To see this, denote by $H[E]$ the linear subspace of \mathbb{R}^{E^*} consisting of all positively homogeneous functions $f : E^* \rightarrow \mathbb{R}$. For $f \in H[E]$ define

$$\|f\|_{\text{FBL}[E]} = \sup \left\{ \sum_{k=1}^n |f(x_k^*)| : n \in \mathbb{N}, x_1^*, \dots, x_n^* \in E^*, \sup_{x \in B_E} \sum_{k=1}^n |x_k^*(x)| \leq 1 \right\}.$$

Given any $x \in E$, let $\delta_x \in H[E]$ be defined by

$$\delta_x(x^*) := x^*(x) \quad \text{for all } x^* \in E^*.$$

It is easy to see that

$$H_1[E] := \{f \in H[E] : \|f\|_{\text{FBL}[E]} < \infty\}$$

is a sublattice of $H[E]$ and that $\|\cdot\|_{\text{FBL}[E]}$ defines a complete lattice norm on $H_1[E]$. Moreover, $\|\delta_x\|_{\text{FBL}[E]} = \|x\|$ for every $x \in E$. As shown in [7], $\text{FBL}[E]$ coincides with the closed sublattice of $H_1[E]$ generated by $\{\delta_x : x \in E\}$ with respect to $\|\cdot\|_{\text{FBL}[E]}$, together with the map $\phi_E(x) = \delta_x$. In particular, by positive homogeneity, we may view elements of $\text{FBL}[E]$ as weak*-continuous functions on the dual ball.

For numerous reasons it is useful to introduce a scale of free Banach lattices, indexed by $p \in [1, \infty]$. Recall that a Banach lattice X is p -convex if there exists a constant $M \geq 1$ such that for every $x_1, \dots, x_m \in X$ we have

$$(1.1) \quad \left\| \left(\sum_{k=1}^m |x_k|^p \right)^{\frac{1}{p}} \right\| \leq M \left(\sum_{k=1}^m \|x_k\|^p \right)^{\frac{1}{p}}.$$

The least constant M satisfying the above inequality is called the p -convexity constant of X and is denoted by $M^{(p)}(X)$. We say that X satisfies an *upper p -estimate* (or that X is a (p, ∞) -convex Banach lattice) if (1.1) holds for all pairwise disjoint vectors $x_1, \dots, x_m \in X$; the least constant is then called the *upper p -estimate constant of X* and is denoted by $M^{(p, \infty)}(X)$. A classical result [22, Proposition 1.f.6] states that a Banach lattice X has an upper p -estimate if and only if

$$(1.2) \quad \left\| \bigvee_{k=1}^m |x_k| \right\| \leq M \left(\sum_{k=1}^m \|x_k\|^p \right)^{\frac{1}{p}}$$

for all $x_1, \dots, x_m \in X$. Moreover, the best choice of M in (1.2) is exactly $M^{(p, \infty)}(X)$. By [11, 22, 30], every p -convex Banach lattice (resp. Banach lattice with an upper p -estimate) can be renormed to have p -convexity (resp. upper p -estimate) constant one.

Given $p \in [1, \infty]$ and a Banach space E , one defines the free p -convex Banach lattice generated by E analogously to $\text{FBL}[E]$: The *free p -convex Banach lattice generated by E* is a Banach lattice $\text{FBL}^{(p)}[E]$ with p -convexity constant one together with a linear isometric embedding $\phi_E : E \rightarrow \text{FBL}^{(p)}[E]$ such that for every p -convex Banach lattice X with p -convexity constant one and every bounded linear operator $T : E \rightarrow X$ there is a unique lattice homomorphism $\hat{T} : \text{FBL}^{(p)}[E] \rightarrow X$ such that $\hat{T} \circ \phi_E = T$ and $\|\hat{T}\| = \|T\|$. By replacing p -convexity with upper p -estimates one obtains the definition of the *free Banach lattice with upper p -estimates generated by E* , which is denoted by $\text{FBL}^{(p, \infty)}[E]$.

As shown in [21], $\text{FBL}^{(p)}[E]$ admits an analogous functional representation to $\text{FBL}[E]$, but with $\|\cdot\|_{\text{FBL}[E]}$ replaced by

$$(1.3) \quad \|f\|_{\text{FBL}^{(p)}[E]} = \sup \left\{ \left(\sum_{k=1}^n |f(x_k^*)|^p \right)^{\frac{1}{p}} : n \in \mathbb{N}, x_1^*, \dots, x_n^* \in E^*, \sup_{x \in B_E} \left(\sum_{k=1}^n |x_k^*(x)|^p \right)^{\frac{1}{p}} \leq 1 \right\}$$

and $H_1[E]$ replaced by

$$H_p[E] := \{f \in H[E] : \|f\|_{\text{FBL}^{(p)}[E]} < \infty\}.$$

Making use of this explicit representation of $\text{FBL}^{(p)}[E]$, most of the major results on $\text{FBL}[E]$ can be shown to hold for $\text{FBL}^{(p)}[E]$. On the other hand, [28, Sections 9.6 and 10] identify several interesting properties of $\text{FBL}^{(p)}[E]$ which depend explicitly on p .

The question of establishing a functional representation for $\text{FBL}^{(p,\infty)}[E]$ was left as an open problem in [21, Remark 6.2] and then reiterated in [28, Section 9.6]. The proof of the existence of $\text{FBL}^{(p,\infty)}[E]$ in [21] proceeds by equipping the free vector lattice with a “maximal” norm satisfying an upper p -estimate and then completing the resulting space. However, this approach gives no practical information about the norm and cannot even guarantee that the resulting Banach lattice is a space of functions on the dual ball. For this reason, minimal progress has been made on $\text{FBL}^{(p,\infty)}[E]$, with the exception of some interesting results being proven in [27].

The first aim of this paper is to answer the above question and identify the norm for $\text{FBL}^{(p,\infty)}[E]$. As a consequence, we obtain the desired continuous injection $\text{FBL}^{(p,\infty)}[E] \hookrightarrow C(B_{E^*})$ and the prospect of ascertaining the fine structure of these spaces. As it turns out, up to equivalence, we have

$$(1.4) \quad \|f\|_{\text{FBL}^{(p,\infty)}[E]} = \sup \left\{ \|(f(x_k^*))_{k=1}^n\|_{\ell^{p,\infty}(n)} : n \in \mathbb{N}, x_1^*, \dots, x_n^* \in E^*, \sup_{x \in B_E} \|(x_k^*(x))_{k=1}^n\|_{\ell^{p,\infty}(n)} \leq 1 \right\},$$

where $\ell^{p,\infty}(n)$ denotes the canonical n -dimensional weak- L^p space. Moreover, as we will see below, the proof that (1.4) is the correct norm is rather illuminating, in that it yields a formula for the norm for the “free Banach lattice” associated to other classes of lattices.

From the above functional description of $\text{FBL}^{(p,\infty)}[E]$, we obtain immediate access to many results. For example, we learn from [6] that $\text{FBL}^{(p,\infty)}[E]$ has only countable disjoint collections and from [7] that lattice homomorphic functionals separate the points of $\text{FBL}^{(p,\infty)}[E]$. With more work, one can prove analogues of many of the results in [28] which were initially proven for $\text{FBL}^{(p)}[E]$. However, certain deeper theorems do not immediately generalize to $\text{FBL}^{(p,\infty)}[E]$. The purpose of the second half of this paper is to prove results of this type. In particular, we solve the *subspace problem* for $\text{FBL}^{(p,\infty)}$ by characterizing when an embedding $\iota : F \hookrightarrow E$ induces a lattice embedding $\bar{\iota} : \text{FBL}^{(p,\infty)}[F] \hookrightarrow \text{FBL}^{(p,\infty)}[E]$. Obtaining such a characterization is by no means trivial, as the proof of the subspace problem for $\text{FBL}^{(p)}$ relied heavily on the L^p -type structure of p -convex Banach lattices.

1.1. Outline of the paper. In Section 2 we study the free Banach lattice generated by E associated to a non-empty class \mathcal{C} of Banach lattices. We begin in Section 2.1 by briefly reviewing the construction of the free vector lattice generated by E . After this, we equip the free vector lattice with the norm

$$\rho_{\mathcal{C}}(f) := \sup \left\{ \|\widehat{T}f\|_X : X \in \mathcal{C}, T : E \rightarrow X \text{ is a linear contraction} \right\},$$

where \widehat{T} denotes the unique lattice homomorphic extension of T , and define the *free Banach lattice generated by E associated to \mathcal{C}* as the completion of $(\text{FVL}[E], \rho_{\mathcal{C}})$. This definition yields a Banach lattice $\text{FBL}^{\mathcal{C}}[E]$ containing an isometric copy of E with the ability to uniquely extend any operator $T : E \rightarrow X \in \mathcal{C}$ to a lattice homomorphism of the same norm. In Section 2.2 we “close” the class \mathcal{C} by constructing a class $\bar{\mathcal{C}} \supseteq \mathcal{C}$ such that $\text{FBL}^{\mathcal{C}}[E] = \text{FBL}^{\bar{\mathcal{C}}}[E] \in \bar{\mathcal{C}}$. This then allows us to identify $\text{FBL}^{\mathcal{C}}[E]$ as a universal object in this new class. In Section 2.3 we find an explicit formula for the norm of $\text{FBL}^{\mathcal{C}}[E]$ when \mathcal{C} consists of a single r.i. space. This, in particular, allows us to identify $\text{FBL}^{\mathcal{C}}[E]$ as a space of weak*-continuous positively homogeneous functions on B_{E^*} . In Section 2.4 we

characterize when continuous injective lattice homomorphisms extend injectively to the completion of a normed lattice, elucidating the difficulty in representing $\text{FBL}^{\mathcal{C}}[E]$ inside of $C(B_{E^*})$.

In Section 3 we combine the results of Section 2 with the Maurey and Pisier factorization theorems to reproduce the norm (1.3) for $\text{FBL}^{(p)}[E]$ and show that (1.4) is a $(1 - \frac{1}{p})^{\frac{1}{p}-1}$ -equivalent lattice norm for $\text{FBL}^{(p,\infty)}[E]$ with upper p -estimate constant one. We also characterize in Theorem 3.8 all of the \mathcal{C} for which we have $\text{FBL}^{\mathcal{C}}[E] = \text{FBL}^{(p)}[E]$ and all of the \mathcal{C}' for which $\text{FBL}^{\mathcal{C}'}[E] \approx \text{FBL}^{(p,\infty)}[E]$. Then, in Section 4 we show that (1.4) is *not* 1-equivalent to the norm satisfying the universal property of $\text{FBL}^{(p,\infty)}[E]$. In fact, by characterizing the sequence spaces that C -lattice embed into $(\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu))_{\infty}$, we are able to show that the entire class of weak- L^p spaces (equipped with any of the renormings in Section 4.1) fails to isometrically witness the norm of the free Banach lattice with upper p -estimates.

Section 5 is devoted to the subspace problem for $\text{FBL}^{(p,\infty)}$; that is, the problem of characterizing those embeddings $\iota : F \hookrightarrow E$ which induce a lattice embedding $\bar{\iota} : \text{FBL}^{(p,\infty)}[F] \hookrightarrow \text{FBL}^{(p,\infty)}[E]$. This is achieved in Section 5.1 by utilizing a novel push-out argument. A benefit of our argument is that it applies equally well to $\text{FBL}^{(p)}$. However, a priori, it yields a slightly different solution to the subspace problem for $\text{FBL}^{(p)}$ than the one in [28, Theorem 3.7]. To reconcile this, we prove the injectivity of ℓ^p in the class of p -convex Banach lattices, which immediately yields the equivalence of the two solutions, up to constants. On the other hand, in Section 5.3 we show that $\ell^{p,\infty}$ is *not* injective in the class of Banach lattices with upper p -estimates, adding an additional novelty to our solution to the subspace problem for $\text{FBL}^{(p,\infty)}[E]$.

2. THE FREE BANACH LATTICE ASSOCIATED TO A CLASS OF BANACH LATTICES

In this section we define and investigate the free Banach lattice $\text{FBL}^{\mathcal{C}}[E]$ generated by a Banach space E associated to a class \mathcal{C} of Banach lattices. We begin in Section 2.1 with a brief review of the construction of the free vector lattice. Then, in Section 2.2, we define $\text{FBL}^{\mathcal{C}}[E]$ as the completion of $\text{FVL}[E]$ under a certain norm, identify an associated class $\bar{\mathcal{C}} \supseteq \mathcal{C}$ of Banach lattices, and prove the universality of $\text{FBL}^{\mathcal{C}}[E]$ in the class $\bar{\mathcal{C}}$. In Section 2.3, an explicit formula for the norm of $\text{FBL}^{\mathcal{C}}[E]$ is obtained when \mathcal{C} consists of a single rearrangement invariant Banach lattice. This, in particular, yields a continuous injection $\text{FBL}^{\mathcal{C}}[E] \hookrightarrow C(B_{E^*})$ for such \mathcal{C} . Finally, in Section 2.4 we supplement the above results with a characterization of the normed lattices for which every continuous lattice homomorphic injection extends injectively to the completion.

2.1. Review of the free vector lattice. Let E be a Banach space and endow B_{E^*} with the weak*-topology. For any $x \in E$, the function $\delta_x : B_{E^*} \rightarrow \mathbb{R}$ given by $\delta_x(x^*) = x^*(x)$ belongs to the space $C(B_{E^*})$ of continuous functions on B_{E^*} . Define a sequence of subspaces of $C(B_{E^*})$ as follows:

$$E_0 = \{\delta_x : x \in E\}; \quad E_n = \text{span}(E_{n-1} \cup \{E_{n-1}\}) \text{ if } n \in \mathbb{N}.$$

Here, for a set A in a vector lattice X , we are denoting $|A| = \{|x| : x \in A\}$. Let $\text{FVL}[E] = \bigcup_{n=0}^{\infty} E_n$ and note that E_0 is a subspace of $C(B_{E^*})$ since $\sum_{k=1}^n a_k \delta_{x_k} = \delta_{\sum_{k=1}^n a_k x_k}$.

Proposition 2.1. *$\text{FVL}[E]$ is the smallest vector sublattice of $C(B_{E^*})$ containing E_0 .*

Proof. Obviously, $\text{span}(E_{n-1}) \subseteq E_n$ for all $n \in \mathbb{N}$. Hence, $\text{FVL}[E]$ is a vector subspace of $C(B_{E^*})$ containing $\{\delta_x : x \in E\}$. Assume that $f \in \text{FVL}[E]$. By definition, $f \in E_n$ for some $n \in \mathbb{N}$. Thus, $|f| \in |E_n| \subseteq E_{n+1} \subseteq \text{FVL}[E]$, so that $\text{FVL}[E]$ is a vector sublattice of $C(B_{E^*})$.

Assume that G is a vector sublattice of $C(B_{E^*})$ containing $\{\delta_x : x \in E\} = E_0$. If $E_{n-1} \subseteq G$ for some $n \in \mathbb{N}$, then $E_{n-1} \cup |E_{n-1}| \subseteq G$. Hence, $E_n \subseteq G$. By induction, $\text{FVL}[E] = \bigcup_{n=0}^{\infty} E_n \subseteq G$. \square

It follows from the expression for $\text{FVL}[E]$ that each $f \in \text{FVL}[E]$ is positively homogeneous in the following sense: $f(\lambda x^*) = \lambda f(x^*)$ if $x^* \in B_{E^*}$ and $0 \leq \lambda \leq 1$. Define $\phi := \phi_E : E \rightarrow \text{FVL}[E]$ by $\phi(x) = \delta_x$. Clearly, ϕ is a linear operator.

Proposition 2.2. *Let $f \in \text{FVL}[E]$. There is a finite subset A of E so that f belongs to the sublattice of $\text{FVL}[E]$ generated by $\{\delta_x : x \in A\}$.*

Proof. The assertion is trivial if $f \in E_0$. Assume that it holds whenever $f \in E_{n-1}$. Let $f \in E_n$ and represent $f = \sum_{k=1}^r (a_k f_k + b_k |f_k|)$ where $a_k, b_k \in \mathbb{R}$ and $f_k \in E_{n-1}$. By the inductive hypothesis, there is a finite set A so that $f_k, 1 \leq k \leq r$, belong to the sublattice generated by $\{\delta_x : x \in A\}$. By construction, f belongs to the same sublattice. \square

An alternative way to prove Propositions 2.1 and 2.2 is to utilize [2, Exercise 8, p. 204]. The following statement is implicit in [28, Theorem 2.1], but we include the details for the convenience of the reader.

Proposition 2.3. *($\text{FVL}[E], \phi$) is the free vector lattice generated by E in the following sense: If X is an Archimedean vector lattice and $T : E \rightarrow X$ is a linear operator then there is a unique linear vector lattice homomorphism $\hat{T} : \text{FVL}[E] \rightarrow X$ such that $\hat{T}\phi = T$.*

Proof. Let A be a finite set in E and let L_A be the sublattice of $C(B_{E^*})$ generated by $\{\delta_x : x \in A\}$. Let $I = I_A$ be the principal lattice ideal in X generated by $\sum_{x \in A} |Tx|$. By standard theory, there is a compact Hausdorff space K and an injective vector lattice homomorphism $i : I \rightarrow C(K)$. For each $t \in K$, define $\ell_t : \text{span } A \rightarrow \mathbb{R}$ by $\ell_t(\sum_{x \in A} a_x x) = \sum_{x \in A} a_x (iT_x)(t)$. Note that if $\sum_{x \in A} a_x x = 0$ then $\sum_{x \in A} a_x Tx = 0$. This implies that $\sum_{x \in A} a_x iT_x = 0$ and thus $\ell_t(\sum_{x \in A} a_x x) = 0$. Hence, ℓ_t is a well-defined linear functional on $\text{span } A$. Since $\text{span } A$ is a finite dimensional subspace of E , ℓ_t is bounded. Let $x_t^* \in E^*$ be a Hahn-Banach extension of ℓ_t .

Suppose that $f \in \text{FVL}[E]$. By Proposition 2.2, there exists a finite set $A = \{x_1, \dots, x_n\}$ in E so that $f \in L_A$. We can therefore write $f = G(\delta_{x_1}, \dots, \delta_{x_n})$ where G is a finite sequence of operations of taking linear combinations and $|\cdot|$. Define $\hat{T}f = G(Tx_1, \dots, Tx_n) \in X$. To see that \hat{T} is well-defined, we have to show that $\hat{T}(\sum a_k f_k) = 0$ if $\sum a_k f_k = 0$, for any finite sum. We may assume that $f_k \in L_A$ for all k . In particular, we have $\hat{T}f_k \in I_A$. Write $f_k = G_k(\delta_{x_1}, \dots, \delta_{x_n})$ for some lattice-linear G_k and note that for any $t \in K$,

$$\begin{aligned} [i\hat{T}(\sum a_k f_k)](t) &= \sum a_k iG_k(Tx_1, \dots, Tx_n)(t) \\ &= \sum a_k G_k(iTx_1(t), \dots, iTx_n(t)) \\ &= \sum a_k G_k(x_t^*(x_1), \dots, x_t^*(x_n)) \\ &= \sum a_k G_k(\delta_{x_1}(x_t^*), \dots, \delta_{x_n}(x_t^*)) \\ &= \sum a_k f_k(x_t^*). \end{aligned}$$

Thus, $\sum a_k f_k = 0$ implies $i\widehat{T}(\sum a_k f_k) = 0$. Since $i : I_A \rightarrow C(K)$ is injective, $\widehat{T}(\sum a_k f_k) = 0$. This proves that \widehat{T} is well-defined. It follows from its definition that \widehat{T} is a linear vector lattice homomorphism.

It is easy to see that for any $x \in E$, $\widehat{T}\phi x = \widehat{T}(\delta_x) = Tx$. Let $S : \text{FVL}[E] \rightarrow X$ be a vector lattice homomorphism so that $S\phi x = Tx$. Then for any $f = G(\delta_{x_1}, \dots, \delta_{x_n})$ we must have

$$Sf = G(S\delta_{x_1}, \dots, S\delta_{x_n}) = G(Tx_1, \dots, Tx_n) = \widehat{T}f.$$

This proves the uniqueness of \widehat{T} . \square

Remark 2.4. In this paper we are identifying $\text{FVL}[E]$ as a space of functions on B_{E^*} . This is in contrast to previous works where $\text{FVL}[E]$ was identified as a space of functions on E^* . By positive homogeneity, these interpretations are clearly equivalent.

2.2. Universality of the free Banach lattice associated to a class of Banach lattices. Let \mathcal{C} be a (non-empty) class of Banach lattices. For instance, \mathcal{C} could be the class of weak- L^p spaces or the class of Banach lattices satisfying an upper p -estimate with constant 1. Given a Banach space E , define a norm $\rho_{\mathcal{C}}$ on $\text{FVL}[E]$ by

$$\rho_{\mathcal{C}}(f) = \sup \left\{ \|\widehat{T}f\|_X : X \in \mathcal{C}, T : E \rightarrow X \text{ is a linear contraction} \right\}.$$

Clearly, $\rho_{\mathcal{C}}$ is a lattice norm on $\text{FVL}[E]$. The *free Banach lattice generated by E associated to \mathcal{C}* is the completion of $(\text{FVL}[E], \rho_{\mathcal{C}})$ and is denoted by $\text{FBL}^{\mathcal{C}}[E]$. It is easy to see that $\phi_E : E \rightarrow \text{FBL}^{\mathcal{C}}[E]$ is a linear isometric embedding. Define an enlarged class of Banach lattices $\overline{\mathcal{C}}$ so that $Y \in \overline{\mathcal{C}}$ if and only if Y is lattice isometric to a closed sublattice of a Banach lattice of the form $(\bigoplus_{\gamma \in \Gamma} X_{\gamma})_{\infty}$, where $X_{\gamma} \in \mathcal{C}$ for all γ . Note that we do not require that the X_{γ} 's be distinct.

Proposition 2.5. *Suppose that $Y \in \overline{\mathcal{C}}$ and $T : E \rightarrow Y$ is a bounded linear operator. Then $\widehat{T} : (\text{FVL}[E], \rho_{\mathcal{C}}) \rightarrow Y$ is a vector lattice homomorphism such that $\|\widehat{T}\| = \|T\|$.*

Proof. Since Y is an Archimedean vector lattice, $\widehat{T} : \text{FVL}[E] \rightarrow Y$ is a vector lattice homomorphism. Note that ϕ_E is an isometric embedding and $\widehat{T}\phi_E = T$. Hence, $\|T\| \leq \|\widehat{T}\|$. On the other hand, suppose that Y is a closed sublattice of X , where $X = (\bigoplus_{\gamma \in \Gamma} X_{\gamma})_{\infty}$, with $X_{\gamma} \in \mathcal{C}$ for all γ . Let π_{γ} be the projection from X onto X_{γ} . Then $\pi_{\gamma}T : E \rightarrow X_{\gamma}$ is a bounded linear operator with $\|\pi_{\gamma}T\| \leq \|T\|$. By definition of $\rho_{\mathcal{C}}$, $\|\widehat{\pi_{\gamma}T}f\| \leq \|T\| \rho_{\mathcal{C}}(f)$ for any $f \in \text{FVL}[E]$. As π_{γ} is a lattice homomorphism, by the uniqueness of $\widehat{\pi_{\gamma}T}$, we have $\widehat{\pi_{\gamma}T} = \pi_{\gamma}\widehat{T}$. Thus, $\|\widehat{T}f\| = \sup_{\gamma} \|\pi_{\gamma}\widehat{T}f\| \leq \|T\| \rho_{\mathcal{C}}(f)$ for any $f \in \text{FVL}[E]$. This shows that $\|\widehat{T}\| \leq \|T\|$. \square

Obviously, the operator \widehat{T} in Proposition 2.5 extends to a lattice homomorphism from $\text{FBL}^{\mathcal{C}}[E]$ to Y , still denoted by \widehat{T} .

Corollary 2.6. *Let E be a Banach space. For any non-empty class \mathcal{C} of Banach lattices, the norms $\rho_{\mathcal{C}}$ and $\rho_{\overline{\mathcal{C}}}$ agree on $\text{FVL}[E]$. In particular, $\text{FBL}^{\mathcal{C}}[E] = \text{FBL}^{\overline{\mathcal{C}}}[E]$ as Banach lattices.*

Proof. Since $\mathcal{C} \subseteq \overline{\mathcal{C}}$ it is clear that $\rho_{\mathcal{C}} \leq \rho_{\overline{\mathcal{C}}}$. On the other hand, suppose that $f \in \text{FVL}[E]$ and $\varepsilon > 0$ are given. There exists $Y \in \overline{\mathcal{C}}$ and a linear contraction $T : E \rightarrow Y$ such that $\rho_{\overline{\mathcal{C}}}(f) \leq (1 + \varepsilon)\|\widehat{T}f\|$. By Proposition 2.5, $\|\widehat{T}f\| \leq \|T\| \rho_{\mathcal{C}}(f) = \rho_{\mathcal{C}}(f)$. It follows that $\rho_{\overline{\mathcal{C}}} \leq \rho_{\mathcal{C}}$. \square

Proposition 2.7. *The Banach lattice $\text{FBL}^{\mathcal{C}}[E]$ belongs to the class $\overline{\mathcal{C}}$.*

Proof. For any $f \in \text{FVL}[E]$ and $n \in \mathbb{N}$, there exists $X(f, n) \in \mathcal{C}$ and a linear contraction $T = T_{f,n} : E \rightarrow X(f, n)$ so that $\rho_{\mathcal{C}}(f) \leq (1 + \frac{1}{n})\|\widehat{T}f\|$. Let $X = (\bigoplus_{f \in \text{FVL}[E], n \in \mathbb{N}} X(f, n))_{\infty}$. Clearly, $X \in \overline{\mathcal{C}}$. Define $j : \text{FBL}^{\mathcal{C}}[E] \rightarrow X$ by $fg = (\widehat{T_{f,n}g})_{f,n}$. Obviously, j is a vector lattice homomorphism. Let $g \in \text{FVL}[E]$. By definition, $\rho_{\mathcal{C}}(g) \geq \|\widehat{T_{f,n}g}\|$ for all (f, n) . Hence, j is a linear contraction. On the other hand, $\|jg\| \geq \|\widehat{T_{g,n}g}\| \geq \frac{n}{n+1}\rho_{\mathcal{C}}(g)$ for all n , implying that $\|jg\| \geq \rho_{\mathcal{C}}(g)$. This shows that $j : \text{FVL}[E] \rightarrow X$ is a lattice isometry. Consequently, $j : \text{FBL}^{\mathcal{C}}[E] \rightarrow X$ is as well. \square

The above discussion motivates the following question.

Problem 2.8. *Find necessary and sufficient conditions on classes of Banach lattices \mathcal{C} and \mathcal{D} so that $\rho_{\mathcal{C}} = \rho_{\mathcal{D}}$, or that $\rho_{\mathcal{C}}$ is equivalent to $\rho_{\mathcal{D}}$.*

In Section 3.1 we will provide a complete solution to Problem 2.8 when \mathcal{C} is the class of p -convex Banach lattices and when \mathcal{C} is the class of Banach lattices with upper p -estimates.

2.3. Rearrangement invariant spaces and the norm for $\text{FBL}^{\mathcal{C}}[E]$. In this subsection, let (Ω, Σ, μ) be a non-atomic σ -finite measure space and let X be a rearrangement invariant (r.i.) Banach function space on Ω in the sense of [9, Definition II.4.1]. To be specific, X is a vector lattice ideal of $L^0(\Omega, \Sigma, \mu)$, equipped with a complete lattice norm $\|\cdot\|$, so that $f \in X, g \in L^0(\Omega, \Sigma, \mu)$ and $\mu\{|f| > t\} = \mu\{|g| > t\}$ for all $t \geq 0$ imply that $g \in X$ and $\|f\| = \|g\|$. We further assume that X has the Fatou property: Whenever $f_n \in X, 0 \leq f_n \uparrow f$ a.e. and $\sup \|f_n\| < \infty$ then $f \in X$ and $\|f\| = \sup_n \|f_n\|$.

Let \mathcal{C} be the class consisting of X only. Given a Banach space E , the aim of this subsection will be to produce a formula for the norm $\rho_{\mathcal{C}}$ on $\text{FVL}[E]$ and show that $\text{FBL}^{\mathcal{C}}[E] \subseteq C(B_{E^*})$.

Given a sequence $\mathcal{U} = (U_n)$ of disjoint measurable subsets of Ω so that $0 < \mu(U_n) < \infty$ for all n , we define for each $f \in L^0(\Omega, \Sigma, \mu)$ the conditional expectation over \mathcal{U} by

$$P_{\mathcal{U}}f = \sum_n \frac{\int_{U_n} f d\mu}{\mu(U_n)} \chi_{U_n}.$$

Note that the above sum has at most one non-zero term at any given point of Ω , so is well-defined as long as $\int_{U_n} f d\mu$ is finite for every $n \in \mathbb{N}$. This will be the case when $f \in X$.

Proposition 2.9. *Let $\mathcal{U} = (U_n)$ be a sequence of disjoint measurable subsets of Ω so that $\mu(U_n) < \infty$ for all n . Then $P_{\mathcal{U}}f \in X$ if $f \in X$ and $P_{\mathcal{U}}$ is a positive contraction on X .*

Proof. First note that, since X is a Banach function space, $\int_{U_n} f d\mu$ is finite for every $n \in \mathbb{N}$ (see property 5 in [9, Definition I.1.1]), so $P_{\mathcal{U}}f$ is well-defined. It is straightforward to check that $P_{\mathcal{U}}$ is a positive contraction from $L^1(\Omega, \Sigma, \mu)$ to $L^1(\Omega, \Sigma, \mu)$ and from $L^{\infty}(\Omega, \Sigma, \mu)$ to $L^{\infty}(\Omega, \Sigma, \mu)$. Thus, $P_{\mathcal{U}}$ is an admissible operator for the compatible pair $(L^1(\Omega, \Sigma, \mu), L^{\infty}(\Omega, \Sigma, \mu))$. Moreover, since X is a r.i. space over a non-atomic σ -finite (in particular, resonant) measure space, it follows from [9, Theorem II.2.2] that X is an exact interpolation space, so $P_{\mathcal{U}} : X \rightarrow X$ is a positive contraction. \square

Lemma 2.10. *Let $h_1, \dots, h_n \in X$ and set $h = \sum_{k=1}^n |h_k|$. Given $\varepsilon > 0$, there is a sequence $\mathcal{U} = (U_n)$ of disjoint measurable subsets of Ω so that $0 < \mu(U_n) < \infty$ for all n and $|h_k - P_{\mathcal{U}}h_k| \leq \varepsilon h, 1 \leq k \leq n$.*

Proof. We may assume that $\varepsilon < 1$. Let $V_i = \{\varepsilon^i \leq h < \varepsilon^{i-1}\}$. Then $(V_i)_{i \in \mathbb{Z}}$ is a sequence of disjoint measurable subsets of Ω so that $h = h_k = 0$ on $(\bigcup_i V_i)^c$ for any k . Since the measure space is σ -finite,

for each i there is an at most countable measurable partition $(U_{ij})_{j \in J_i}$ of V_i with $0 < \mu(U_{ij}) < \infty$ and $b_{kij} \in \mathbb{R}$ so that

$$|h_k \chi_{V_i} - \sum_{j \in J_i} b_{kij} \chi_{U_{ij}}| \leq \frac{\varepsilon^{i+1}}{2}, \quad i \in \mathbb{Z}, \quad 1 \leq k \leq n.$$

Thus, for each $1 \leq k \leq n$ we have

$$(2.1) \quad |h_k - \sum_{i,j} b_{kij} \chi_{U_{ij}}| \leq \sum_i \frac{\varepsilon^{i+1}}{2} \chi_{V_i} \leq \frac{\varepsilon h}{2}$$

and

$$\left| \frac{1}{\mu(U_{ij})} \int_{U_{ij}} h_k d\mu - b_{kij} \right| = \left| \frac{1}{\mu(U_{ij})} \int_{U_{ij}} h_k - b_{kij} \chi_{U_{ij}} d\mu \right| \leq \frac{\varepsilon^{i+1}}{2}.$$

Take $\mathcal{U} = (U_{ij})_{i \in \mathbb{Z}, j \in J_i}$. Note that $\mathcal{P}_{\mathcal{U}} h_k = 0$ on the set $(\bigcup_{ij} U_{ij})^c$. For $1 \leq k \leq n$,

$$(2.2) \quad |\mathcal{P}_{\mathcal{U}} h_k - \sum_{i,j} b_{kij} \chi_{U_{ij}}| \leq \sum_{i,j} \frac{\varepsilon^{i+1}}{2} \chi_{U_{ij}} = \sum_i \frac{\varepsilon^{i+1}}{2} \chi_{V_i} \leq \frac{\varepsilon h}{2}.$$

Summing (2.1) and (2.2) gives the desired result. \square

Let $I = \{(m, r) \in \mathbb{N}^2 : \frac{m}{r} \leq \mu(\Omega)\}$. If $(m, r) \in I$, take any sequence $(V_i)_{i=1}^m$ of disjoint measurable subsets of Ω such that $\mu(V_i) = r^{-1}$, $1 \leq i \leq m$. Define a norm ρ_{mr} on \mathbb{R}^m by

$$\rho_{mr}(a_1, \dots, a_m) = \left\| \sum_{i=1}^m a_i \chi_{V_i} \right\|_X.$$

Let E be a Banach space and define $\rho : C(B_{E^*}) \rightarrow [0, \infty]$ by

$$\rho(f) = \sup \rho_{mr}(f(x_1^*), \dots, f(x_m^*)),$$

where the supremum is taken over all $(m, r) \in I$ and $x_1^*, \dots, x_m^* \in E^*$ which satisfy the constraint $\sup_{x \in B_E} \rho_{mr}(x_1^*(x), \dots, x_m^*(x)) \leq 1$. By definition, $\rho(\delta_x) = \|x\|$ for any $x \in E$. Moreover, it is easy to check that ρ is a complete lattice norm on the vector lattice $\{f \in C(B_{E^*}) : \rho(f) < \infty\}$ which contains $\text{FVL}[E]$. For our next result, recall that we are considering the case $\mathcal{C} = \{X\}$.

Theorem 2.11. *The norms ρ and $\rho_{\mathcal{C}}$ are equal on $\text{FVL}[E]$. In particular, $\text{FBL}^{\mathcal{C}}[E] \subseteq \{f \in C(B_{E^*}) : \rho(f) < \infty\}$.*

Proof. Suppose that $f \in \text{FVL}[E]$. There are $x_1, \dots, x_n \in E$ and a lattice-linear function G so that $f = G(\delta_{x_1}, \dots, \delta_{x_n})$. Let $\varepsilon > 0$ be given. There are $(m, r) \in I$ and $x_1^*, \dots, x_m^* \in E^*$ so that $\sup_{x \in B_E} \rho_{mr}(x_1^*(x), \dots, x_m^*(x)) \leq 1$ and

$$\rho(f) \leq \rho_{mr}(f(x_1^*), \dots, f(x_m^*)) + \varepsilon = \left\| \sum_{i=1}^m f(x_i^*) \chi_{V_i} \right\|_X + \varepsilon,$$

where $(V_i)_{i=1}^m$ is the sequence of disjoint measurable subsets of Ω chosen above, so that $\mu(V_i) = r^{-1}$, $1 \leq i \leq m$. Define a linear operator $S : \text{FVL}[E] \rightarrow X$ by $Sg = \sum_{i=1}^m g(x_i^*) \chi_{V_i}$. It is clear that S is a lattice homomorphism. By definition of ρ , if $\rho(g) \leq 1$, then $\rho_{mr}(g(x_1^*), \dots, g(x_m^*)) \leq 1$ and thus $\|Sg\| \leq 1$. Let $T = S\phi_E : E \rightarrow X$. Then $\|T\| \leq 1$ and hence

$$\rho_{\mathcal{C}}(f) \geq \|\widehat{T}f\| = \|\widehat{S\phi_E}f\| = \|G(S\delta_{x_1}, \dots, S\delta_{x_n})\| = \|Sf\|,$$

where the last step holds since S is a lattice homomorphism. Thus, $\rho_C(f) \geq \rho(f) - \varepsilon$ for any $\varepsilon > 0$. Therefore, $\rho_C(f) \geq \rho(f)$.

Conversely, given $\varepsilon > 0$, there exists a linear contraction $T : E \rightarrow X$ so that $\rho_C(f) \leq \|\widehat{T}f\| + \varepsilon$. Let $h_k = Tx_k$, $1 \leq k \leq n$, and $h = \sum_{k=1}^n |h_k|$. By Lemma 2.10, there is a sequence $\mathcal{U} = (U_i)$ of disjoint measurable subsets of Ω so that $0 < \mu(U_i) < \infty$ for all i and $|h_k - \mathcal{P}_U h_k| \leq \varepsilon h$, $1 \leq k \leq n$. There is a constant $C_G < \infty$, depending only on G , so that

$$|\widehat{T}f - G(\mathcal{P}_U h_1, \dots, \mathcal{P}_U h_n)| = |G(h_1, \dots, h_n) - G(\mathcal{P}_U h_1, \dots, \mathcal{P}_U h_n)| \leq C_G \varepsilon h.$$

Express $\mathcal{P}_U h_k$ as $\sum_i a_{ki} \chi_{U_i}$, where $T^* \chi_{U_i} = y_i^*$ and

$$a_{ki} = \frac{1}{\mu(U_i)} \int_{U_i} h_k d\mu = \frac{y_i^*(x_k)}{\mu(U_i)}.$$

Then

$$\begin{aligned} G(\mathcal{P}_U h_1, \dots, \mathcal{P}_U h_n) &= G\left(\sum_i a_{1i} \chi_{U_i}, \dots, \sum_i a_{ni} \chi_{U_i}\right) \\ &= \sum_i G(a_{1i}, \dots, a_{ni}) \chi_{U_i} \\ &= \sum_i G(\delta_{x_1}, \dots, \delta_{x_n})(y_i^*) \frac{\chi_{U_i}}{\mu(U_i)} \\ &= \sum_i \frac{f(y_i^*)}{\mu(U_i)} \chi_{U_i}. \end{aligned}$$

Since X has the Fatou property, there exists $l \in \mathbb{N}$ so that

$$\left\| \sum_{i=1}^l \frac{f(y_i^*)}{\mu(U_i)} \chi_{U_i} \right\| > \|G(\mathcal{P}_U h_1, \dots, \mathcal{P}_U h_n)\| - \varepsilon \geq \|\widehat{T}f\| - (1 + C_G \|h\|) \varepsilon.$$

Using the Fatou property once more, choose disjoint Lebesgue measurable sets V_1, \dots, V_l so that $V_i \subseteq U_i$, $\mu(V_i) \in \mathbb{Q}$ and

$$\|\widehat{T}f\| - (1 + C_G \|h\|) \varepsilon < \left\| \sum_{j=1}^l \frac{f(y_j^*)}{\mu(U_j)} \chi_{V_j} \right\|.$$

Write $\mu(V_i) = \frac{j_i}{r}$, where $j_i, r \in \mathbb{N}$. Decompose each V_i as a disjoint union $V_i = \bigcup_{s=1}^{j_i} V_{is}$, with $\mu(V_{is}) = r^{-1}$. Take $j_0 = \sum_{i=1}^l j_i$. It follows that $(j_0, r) \in I$. Consider the sequence where each $\frac{y_i^*}{\mu(U_i)}$ is repeated j_i times. For any $x \in B_E$, let b be the element of \mathbb{R}^{j_0} so that each $\frac{y_i^*(x)}{\mu(U_i)}$ occurs j_i times. Then

$$\rho_{j_0 r}(b) = \left\| \sum_{i=1}^l \frac{y_i^*(x)}{\mu(U_i)} \chi_{V_i} \right\| \leq \|\mathcal{P}_U(Tx)\| \leq \|T\| = 1.$$

Recall that elements of $\text{FVL}[E]$ have a homogeneity property given after Proposition 2.1. It follows from the definition of ρ that, taking a to be the element of \mathbb{R}^{j_0} where each $\frac{f(y_i^*)}{\mu(U_i)}$ occurs j_i times,

$$\begin{aligned} \rho(f) &\geq \rho_{j_0 r}(a) = \left\| \sum_{i=1}^l \sum_{s=1}^{j_i} \frac{f(y_i^*)}{\mu(U_i)} \chi_{V_{is}} \right\| = \left\| \sum_{i=1}^l \frac{f(y_i^*)}{\mu(U_i)} \chi_{V_i} \right\| \\ &> \|\widehat{T}f\| - (1 + C_G \|h\|) \varepsilon \geq \rho_C(f) - (2 + C_G \|h\|) \varepsilon. \end{aligned}$$

Thus, $\rho(f) \geq \rho_C(f)$. \square

The above construction of $\text{FBL}^C[E]$ motivates the following question.

Question 2.12. It is well-known that the properties of $\text{FBL}^{(p)}[E]$ depend heavily on whether p is finite or infinite. Can the Banach lattices $\text{FBL}^C[E]$ be used to interpolate such properties? For example, if (e_k) is the unit vector basis of ℓ^2 , then by [28, Propositions 5.14 and 6.4] $(|\delta_{e_k}|)$ gives a copy of ℓ^1 in $\text{FBL}^{(p)}[\ell^2]$ when $p < \infty$ and a copy of ℓ^2 when $p = \infty$. Can we find for each $r \in (1, 2)$ a class \mathcal{C}_r so that $(|\delta_{e_k}|)$ behaves like ℓ^r in $\text{FBL}^{\mathcal{C}_r}[\ell^2]$?

2.4. Injective homomorphisms need not extend to the norm completion. In Section 2.2 we defined $\text{FBL}^C[E]$ as the completion of $(\text{FVL}[E], \rho_C)$. Clearly, there is a continuous lattice homomorphic injection $(\text{FVL}[E], \rho_C) \hookrightarrow C(B_{E^*})$ because of the inequality $\|\cdot\|_\infty \leq \rho_C(\cdot)$ on $\text{FVL}[E]$. In [13] it is shown that under certain order continuity assumptions (which are not fulfilled in our situation) continuous lattice homomorphic injections retain their injectivity after norm completion. Moreover, it is claimed in [39, Lemma 14] that *any* continuous lattice homomorphic injection from a normed vector lattice to a Banach lattice lifts injectively to the completion. If this were the case, it would provide a straightforward solution to the question of finding a function lattice representation for $\text{FBL}^{(p, \infty)}[E]$ asked in [21, Remark 6.2]. In this subsection, we give a complete characterization of when continuous injective lattice homomorphisms extend injectively to the completion, and, in particular, show that this need not always be the case.

Throughout this subsection X will be a vector lattice equipped with a lattice norm and Y will be a Banach lattice. We denote the norm completion of X by \widehat{X} and consider polar sets with respect to the duality $\langle \widehat{X}, X^* \rangle$: for $A \subseteq \widehat{X}$, set $A^\circ = \{x^* \in X^* : \langle a, x^* \rangle \leq 1, \forall a \in A\}$.

Theorem 2.13. *Let X be a normed vector lattice equipped with a lattice norm and let \widehat{X} be its norm completion. The following are equivalent.*

- (1) *For any Banach lattice Y , any bounded linear injective lattice homomorphism $T : X \rightarrow Y$ extends to a bounded linear injective operator $\widehat{T} : \widehat{X} \rightarrow Y$.*
- (2) *For any $0 < x \in \widehat{X}$, $I_x \cap X \neq \{0\}$, where I_x is the closed ideal in \widehat{X} generated by x .*

Proof. (1) \implies (2): Suppose that there exists $0 < x \in \widehat{X}$ so that $I_x \cap X = \{0\}$. By [38, Proposition II.4.7], $(I_x)^\circ$ is an ideal in X^* . Define $\rho : X \rightarrow \mathbb{R}$ by

$$\rho(u) = \sup_{x^* \in (I_x)^\circ \cap B_{X^*}} |\langle u, x^* \rangle|.$$

Since $(I_x)^\circ \cap B_{X^*}$ is a solid subset of B_{X^*} , ρ is a lattice semi-norm on X such that $\rho(\cdot) \leq \|\cdot\|$. If $0 < u \in X$, then $u \notin I_x$. Since I_x is a closed subspace of \widehat{X} , there exists $x^* \in (I_x)^\circ \cap B_{X^*}$ such that $\langle u, x^* \rangle \neq 0$. Hence, $\rho(u) > 0$. This shows that ρ is a norm on X .

Denote by X_ρ the vector lattice X normed by ρ and let $i : X \rightarrow X_\rho \subseteq \widehat{X}_\rho$ be the formal identity, where \widehat{X}_ρ is the norm completion of X_ρ . By construction, i is a bounded linear injective lattice homomorphism. By (1), i extends to an injective bounded linear operator $\widehat{i} : \widehat{X} \rightarrow \widehat{X}_\rho$. Denote by $\widehat{\rho}$ the norm on \widehat{X}_ρ . By injectivity of \widehat{i} , $\widehat{i}x \neq 0$. Let $c = \widehat{\rho}(\widehat{i}x) > 0$. There exists (x_n) in X that converges to x in \widehat{X} . By continuity, (ix_n) converges in \widehat{X}_ρ to $\widehat{i}x$. Choose N large enough so that $\widehat{\rho}(ix_N - \widehat{i}x) < \frac{c}{2}$. Then $\rho(ix_N) > \frac{c}{2}$. By definition of ρ , there exists $x^* \in (I_x)^\circ \cap B_{X^*}$ so that

$|\langle ix_N, x^* \rangle| > \frac{c}{2}$. By definition of ρ again, x^* defines a bounded linear functional on X_ρ with norm $\rho^*(x^*) \leq 1$. Hence,

$$|\langle x, x^* \rangle| = |\langle \widehat{ix}, x^* \rangle| \geq |\langle ix_N, x^* \rangle| - \widehat{\rho}(ix_N - \widehat{ix}) \cdot \rho^*(x^*) > \frac{c}{2} - \frac{c}{2} = 0.$$

This is impossible since $x \in I_x$ and $x^* \in (I_x)^\circ$.

(2) \implies (1): Let $T : X \rightarrow Y$ be as in (1), and assume that its extension \widehat{T} is not injective. There exists some $x \in \widehat{X}$, $x \neq 0$, such that $\widehat{T}x = 0$. Since \widehat{T} is a lattice homomorphism, we can assume that $x > 0$. Let I_x be the closed ideal in \widehat{X} generated by x . By (2), there is a non-zero element $z \in I_x \cap X$. Since I_x is the closure of the ideal generated by x , there exists a sequence $(z_n) \subseteq \widehat{X}$ converging to z in \widehat{X} and some positive scalars (λ_n) such that $0 \leq |z_n| \leq \lambda_n x$. Therefore,

$$0 \leq |\widehat{T}z_n| \leq \lambda_n \widehat{T}x = 0$$

and

$$Tz = \widehat{T}z = \lim_n \widehat{T}z_n = 0,$$

so by the injectivity of T we conclude that z must be zero. This is a contradiction. \square

The following example taken from [10] illustrates the above situation.

Example 2.14. *There is a dense vector sublattice X of c_0 such that $I \cap X = \{0\}$, where I is the closed ideal in c_0 generated by e_1 ; that is, $I = \{(a_n) \in c_0 : a_n = 0 \text{ for all } n \geq 2\}$. Consequently, not every bounded linear injective lattice homomorphism from X into a Banach lattice Y extends to an injective operator from c_0 to Y .*

Proof. Let X consist of all $(a_n) \in c_0$ such that there exists $m \in \mathbb{N}$ so that $a_n = \frac{a_1}{n}$ for all $n \geq m$. Clearly, X is a vector sublattice of c_0 with $e_n \in X$ if $n \geq 2$. For any $n \geq 2$, $x_n := e_1 + \sum_{k=n}^{\infty} \frac{e_k}{k} \in X$ and (x_n) converges to e_1 in c_0 . Thus, $e_1 \in \overline{X}$. It follows that $\overline{X} = c_0$. If $a = (a_n) \in I \cap X$, then $a_n = 0$ for all $n \geq 2$. Since $a_1 = ma_m$ for sufficiently large m , $a_1 = 0$ as well. Thus, $a = 0$. \square

3. APPLICATIONS TO FREE p -CONVEX AND FREE (p, ∞) -CONVEX BANACH LATTICES

We now apply the results of Sections 2.2 and 2.3 to the classes \mathcal{C}_p and $\mathcal{C}_{p,\infty}$ consisting of all Banach lattices with p -convexity constant 1 and all Banach lattices with upper p -estimate constant 1, respectively. For this purpose, we recall the following factorization theorems.

Theorem 3.1 (Maurey's Factorization Theorem). *Let $1 < p < \infty$, $A \subseteq L^1(\mu)$ and $0 < M < \infty$. The following are equivalent.*

(1) *For all finitely supported sequences $(\alpha_i)_{i \in I}$ of real numbers and $(f_i)_{i \in I} \subseteq A$,*

$$\left\| \left(\sum |\alpha_i f_i|^p \right)^{\frac{1}{p}} \right\|_{L^1(\mu)} \leq M \left(\sum |\alpha_i|^p \right)^{\frac{1}{p}}.$$

(2) *There exists $g \in L^1(\mu)_+$, $\|g\|_1 = 1$, such that for any $f \in A$,*

$$\left\| \frac{f}{g} \right\|_{L^p(g \cdot \mu)} \leq M.$$

Theorem 3.2 (Pisier's Factorization Theorem). *Let $1 < p < \infty$, $A \subseteq L^1(\mu)$, $0 < M < \infty$ and $\gamma_p = (1 - \frac{1}{p})^{\frac{1}{p}-1}$. Consider the following conditions.*

(1) For all finitely supported sequences $(\alpha_i)_{i \in I}$ of real numbers and $(f_i)_{i \in I} \subseteq A$,

$$\left\| \bigvee |\alpha_i f_i| \right\|_{L^1(\mu)} \leq M \left(\sum |\alpha_i|^p \right)^{\frac{1}{p}}.$$

(2) There exists $g \in L^1(\mu)_+$, $\|g\|_1 \leq 1$, such that for any $f \in A$ and any μ -measurable set U ,

$$\|f \chi_U\|_{L^1(\mu)} \leq \gamma_p M \left(\int_U g d\mu \right)^{1-\frac{1}{p}}.$$

Then (1) \implies (2).

Theorem 3.1 is very well-known. See, for example, [1, Chapter 7]. Theorem 3.2 is the implication (iii) \implies (i) in [30, Theorem 1.1]. Note that condition (2) in Theorem 3.2 is equivalent to

$$\left\| \frac{f}{g} \chi_U \right\|_{L^1(g \cdot \mu)} \leq \gamma_p M \left((g \cdot \mu)(U) \right)^{1-\frac{1}{p}}.$$

In particular, under the above conditions, $\left\| \frac{f}{g} \right\|_{L^{p,\infty}(g \cdot \mu)} \leq \gamma_p M$ for all $f \in A$.

We will use the above factorization theorems to represent p -convex Banach lattices inside of infinity sums of L^p spaces and Banach lattices with upper p -estimates inside of infinity sums of weak- L^p spaces. Recall that, given a measure space (Ω, Σ, μ) , the space $L^{p,\infty}(\mu)$ is defined as the set of all measurable functions $h : \Omega \rightarrow \mathbb{R}$ such that the quasinorm

$$\sup_{t>0} t \mu(\{|h| > t\})^{\frac{1}{p}}$$

is finite. Note, in particular, that every $h \in L^{p,\infty}(\mu)$ has σ -finite support. Unless otherwise specified, we will equip $L^{p,\infty}(\mu)$ with the norm

$$\|h\|_{L^{p,\infty}} = \sup \left\{ \mu(E)^{\frac{1}{p}-1} \int_E |h| d\mu : 0 < \mu(E) < \infty \right\}.$$

It is a standard fact (cf. [19, Exercise 1.1.12]) that the above norm is equivalent to the usual quasinorm on weak- L^p ; it can also be checked to be a lattice norm with upper- p estimate constant one.

Proposition 3.3. *The following statements hold:*

(1) Let X be a p -convex Banach lattice with constant M . There is a family Γ of probability measures and a lattice isomorphism

$$J : X \rightarrow \left(\bigoplus_{\mu \in \Gamma} L^p(\mu) \right)_{\infty}$$

such that $\|x\| \leq \|Jx\| \leq M\|x\|$ for all $x \in X$.

(2) Let X be a (p, ∞) -convex Banach lattice with constant M . There is a family Γ of probability measures and a lattice isomorphism

$$J : X \rightarrow \left(\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu) \right)_{\infty}$$

such that $\|x\| \leq \|Jx\| \leq \gamma_p M \|x\|$ for all $x \in X$.

Proof. Let $x \in S_X^+ = \{z \in X_+ : \|z\| = 1\}$. Choose $x^* \in X_+^*$, $\|x^*\| = 1$, such that $x^*(x) = 1$. Define $\rho_x : X \rightarrow \mathbb{R}$ by $\rho_x(z) = x^*(|z|)$. Then ρ_x is an L -norm on $X/\ker \rho_x$. Let $q_x : X \rightarrow X/\ker \rho_x$ be the quotient map. There exist a probability measure μ_x and a contractive lattice homomorphism $i_x : X/\ker \rho_x \rightarrow L^1(\mu_x)$ such that $i_x q_x x = 1$, the constant 1 function. Set $A_x = i_x q_x(B_X) \subseteq L^1(\mu_x)$. Let α_i be a finitely supported real sequence and let $f_i = i_x q_x x_i$ for $x_i \in B_X$. Since $i_x q_x$ is a lattice homomorphism,

$$\begin{aligned} \left\| \left(\sum |\alpha_i f_i|^p \right)^{\frac{1}{p}} \right\|_{L^1(\mu_x)} &= \left\langle \left(\sum |\alpha_i x_i|^p \right)^{\frac{1}{p}}, x^* \right\rangle \leq \left\| \left(\sum |\alpha_i x_i|^p \right)^{\frac{1}{p}} \right\| \\ &\leq M \left(\sum |\alpha_i|^p \right)^{\frac{1}{p}} \text{ if } X \text{ is } p\text{-convex with constant } M, \\ \left\| \bigvee |\alpha_i f_i| \right\|_{L^1(\mu_x)} &= \left\langle \bigvee |\alpha_i x_i|, x^* \right\rangle \leq \left\| \bigvee |\alpha_i x_i| \right\| \\ &\leq M \left(\sum |\alpha_i|^p \right)^{\frac{1}{p}} \text{ if } X \text{ is } (p, \infty)\text{-convex with constant } M. \end{aligned}$$

By the factorization theorems above, there exists $g_x \in L^1(\mu_x)_+$, $\|g_x\|_{L^1(\mu_x)} \leq 1$, such that for any $f \in A$ and any μ_x measurable set U ,

$$\left\| \frac{f}{g_x} \right\|_{L^p(g_x \cdot \mu_x)} \leq M, \text{ respectively, } \left\| \frac{f}{g_x} \right\|_{L^{p,\infty}(g_x \cdot \mu_x)} \leq \gamma_p M.$$

Define

$$J : X \rightarrow \left(\bigoplus_{x \in S_X^+} L^p(g_x \cdot \mu_x) \right)_\infty, \text{ respectively } J : X \rightarrow \left(\bigoplus_{x \in S_X^+} L^{p,\infty}(g_x \cdot \mu_x) \right)_\infty$$

by $Jz = (i_x q_x z / g_x)_x$. It is easy to check that J is a lattice homomorphism. If $z \in B_X$, we have $i_x q_x z \in A_x$. Hence,

$$\|i_x q_x z / g_x\|_{L^p(g_x \cdot \mu_x)} \leq M, \text{ respectively } \|i_x q_x z / g_x\|_{L^{p,\infty}(g_x \cdot \mu_x)} \leq \gamma_p M.$$

Thus, $\|Jz\| \leq M\|z\|$, respectively $\|Jz\| \leq \gamma_p M\|z\|$ for all $z \in X$.

On the other hand, if $x \in S_X^+$, then in the p -convex case,

$$\|Jx\| \geq \|i_x q_x x / g_x\|_{L^p(g_x \cdot \mu_x)} \geq \|i_x q_x x / g_x\|_{L^1(g_x \cdot \mu_x)}$$

since $\|g_x \cdot \mu_x\| \leq 1$. Therefore, $\|Jx\| \geq \|i_x q_x x\|_{L^1(\mu_x)} = x^*(x) = 1$. In the (p, ∞) -convex case, suppose that μ_x is a measure defined on Ω_x . Then

$$\int \frac{i_x q_x x}{g_x} d(g_x \cdot \mu_x) = \int i_x q_x x d\mu_x = x^*(x) = 1 \geq \left((g_x \cdot \mu_x)(\Omega_x) \right)^{1-\frac{1}{p}}.$$

Hence, $\|Jx\| \geq \left\| \frac{i_x q_x x}{g_x} \right\|_{L^{p,\infty}(g_x \cdot \mu_x)} \geq 1$. It follows that $\|Jx\| \geq \|x\|$ for all $x \in X$ in both cases. \square

Proposition 3.3 has the following interpretation. Assume that $1 < p < \infty$. Let \mathcal{C}_p be the class of all p -convex Banach lattices with constant 1 and let \mathcal{X}_p be the class of all $L^p(\mu)$ spaces for any measure μ . Then $\mathcal{C}_p \subseteq \overline{\mathcal{X}_p}$. Since the reverse inclusion is evident, we have $\mathcal{C}_p = \overline{\mathcal{X}_p}$. By Corollary 2.6, for any Banach space E , $\rho_{\mathcal{C}_p} = \rho_{\mathcal{X}_p}$ on $\text{FVL}[E]$. Note further that, since the measures $\mu \in \Gamma$ in Proposition 3.3 are finite, we could restrict the class \mathcal{X}_p to the class of $L^p(\mu)$ spaces with finite measures (let us denote this class by \mathcal{X}_p^F) and the equality of norms would still hold. Actually, it can be proved by a standard argument that $\mathcal{X}_p^F \subseteq \mathcal{X}_p \subseteq \overline{\mathcal{X}_p^F}$. The following result gives an alternative method for computing $\rho_{\mathcal{C}_p}$.

Theorem 3.4. *Let E be a Banach space and let $1 < p < \infty$. The norm $\rho_{\mathcal{C}_p}$ on $\text{FVL}[E]$ is given by*

$$\rho_{\mathcal{C}_p}(f) = \rho_p(f) := \sup \|(f(x_1^*), \dots, f(x_n^*))\|_p,$$

where the supremum is taken over all finite collections $x_1^*, \dots, x_n^* \in B_{E^*}$ satisfying the constraint $\sup_{x \in B_E} \|(x_1^*(x), \dots, x_n^*(x))\|_p \leq 1$.

Proof. As observed above, $\rho_{\mathcal{C}_p} = \rho_{\mathcal{X}_p} = \rho_{\mathcal{X}_p^F}$, where \mathcal{X}_p^F is the class of all $L^p(\mu)$ spaces with finite μ . If $X \in \mathcal{X}_p^F$ there exists a non-atomic finite measure μ so that X is lattice isometric to a sublattice of $L^p(\mu)$. Thus, $\rho_{\{X\}} \leq \rho_{\{L^p(\mu)\}}$, where the two norms here are the ones associated to the classes with single elements, as indicated. By Theorem 2.11, $\rho_{\{L^p(\mu)\}}(f) = \rho_p(f)$. Finally, note that

$$\rho_{\mathcal{X}_p^F} = \sup_{X \in \mathcal{X}_p^F} \rho_{\{X\}} \leq \sup_{\substack{\mu \text{ non-atomic} \\ \text{finite}}} \rho_{\{L^p(\mu)\}} = \rho_p \leq \rho_{\mathcal{X}_p^F}.$$

Hence, $\rho_{\mathcal{C}_p} = \rho_p$, as claimed. \square

Let $\mathcal{C}_{p,\infty}$ be the class of all (p, ∞) -convex Banach lattices with constant 1 and let $\mathcal{X}_{p,\infty}^\sigma$ be the class of all $L^{p,\infty}(\mu)$ spaces for any σ -finite measure μ . Similarly to Theorem 3.4, we have the following result.

Theorem 3.5. *Let E be a Banach space and let $1 < p < \infty$. The norm $\rho_{\mathcal{X}_{p,\infty}^\sigma}$ on $\text{FVL}[E]$ is given by*

$$\rho_{\mathcal{X}_{p,\infty}^\sigma}(f) = \rho_{p,\infty}(f) := \sup \|(f(x_1^*), \dots, f(x_n^*))\|_{p,\infty},$$

where the supremum is taken over all finite collections $x_1^*, \dots, x_n^* \in B_{E^*}$ satisfying the constraint $\sup_{x \in B_E} \|(x_1^*(x), \dots, x_n^*(x))\|_{p,\infty} \leq 1$. Moreover, $\rho_{p,\infty} \leq \rho_{\mathcal{C}_{p,\infty}} \leq \gamma_p \rho_{p,\infty}$, where $\gamma_p = (1 - \frac{1}{p})^{\frac{1}{p}-1}$.

Proof. The proof that $\rho_{\mathcal{X}_{p,\infty}^\sigma} = \rho_{p,\infty}$ is similar to the proof of Theorem 3.4. Since $\mathcal{X}_{p,\infty}^\sigma \subseteq \mathcal{C}_{p,\infty}$, $\rho_{\mathcal{X}_{p,\infty}^\sigma} \leq \rho_{\mathcal{C}_{p,\infty}}$. On the other hand, let $X \in \mathcal{C}_{p,\infty}$. By Proposition 3.3, there exists $Y \in \overline{\mathcal{X}_{p,\infty}^\sigma}$ and a lattice isomorphism $J : X \rightarrow Y$ so that $\|x\| \leq \|Jx\| \leq \gamma_p \|x\|$ for all $x \in X$. Let $T : E \rightarrow X$ be a linear contraction. Then $\gamma_p^{-1}JT : E \rightarrow Y$ is a linear contraction. Since J is a lattice homomorphism, $\widehat{\gamma_p^{-1}JT} = \gamma_p^{-1}J\widehat{T}$. For any $f \in \text{FVL}[E]$,

$$\|\widehat{T}f\| \leq \gamma_p \|\gamma_p^{-1}J\widehat{T}f\| = \gamma_p \|\widehat{\gamma_p^{-1}JT}f\| \leq \gamma_p \rho_{\mathcal{X}_{p,\infty}^\sigma}(f).$$

Taking supremum over all linear contractions $T : E \rightarrow X$ for any $X \in \mathcal{C}_{p,\infty}$ shows that $\rho_{\mathcal{C}_{p,\infty}} \leq \gamma_p \rho_{\mathcal{X}_{p,\infty}^\sigma}$. \square

Observe that Theorems 3.4 and 3.5 say that $\rho_{\mathcal{C}_p} = \rho_{\{\ell^p\}}$ and that $\rho_{\{\ell^{p,\infty}\}} \leq \rho_{\mathcal{C}_{p,\infty}} \leq \gamma_p \rho_{\{\ell^{p,\infty}\}}$. As a result, $\text{FBL}^{\mathcal{C}_p}[E] = \text{FBL}^{\{\ell^p\}}[E]$ and $\text{FBL}^{\mathcal{C}_{p,\infty}}[E] = \text{FBL}^{\{\ell^{p,\infty}\}}[E]$ and it is clear that both of these spaces are vector sublattices of $C(B_{E^*})$. The proof works in exactly the same way if $p = 1$, in which case we get that \mathcal{C}_1 , the class of all (1-convex) Banach lattices, is equal to $\overline{\mathcal{X}_1}$, where \mathcal{X}_1 consists of all $L^1(\mu)$ spaces. The Banach lattice $\text{FBL}^{\mathcal{C}_1}[E]$ is the free Banach lattice generated by E , which is usually denoted by $\text{FBL}[E]$. From the above, we see that $\rho_{\mathcal{C}_1} = \rho_{\mathcal{X}_1} = \rho_{\{\ell^1\}}$.

Remark 3.6. In this subsection we have made heavy use of factorization theorems through L^p and weak- L^p to identify the norm of the associated free spaces as a type of nonlinear weak-strong summing norm. We note that there is a large literature (see [24, 25, 26, 36] and related papers of these authors) on factorization theory and generalized summing operators on Banach function

spaces, which may prove to be useful in the study of $\text{FBL}^{\mathcal{C}}[E]$. There are also several papers [16, 32, 33, 34, 35, 37] which study factorizations of p -convex and q -concave operators. Although we will not do so here, it is certainly possible to extend some of these results to other convexity/concavity conditions, such as (p, ∞) -convexity and $(q, 1)$ -concavity.

3.1. Solution to Problem 2.8 for the classes \mathcal{C}_p and $\mathcal{C}_{p,\infty}$. Let \mathcal{C}_p and $\mathcal{C}_{p,\infty}$ be the class of p -convex Banach lattices and the class of Banach lattices satisfying an upper p -estimate with constant 1, respectively. We will characterize the classes of Banach lattices \mathcal{D} so that $(\text{FBL}^{\mathcal{C}_p}[E], \rho_{\mathcal{C}_p})$ and $(\text{FBL}^{\mathcal{D}}[E], \rho_{\mathcal{D}})$ agree for any Banach space E and also the classes \mathcal{D}' so that $(\text{FBL}^{\mathcal{C}_{p,\infty}}[E], \rho_{\mathcal{C}_{p,\infty}})$ and $(\text{FBL}^{\mathcal{D}'}[E], \rho_{\mathcal{D}'})$ are lattice isomorphic.

For the proofs of the above claims, we need an extension of the isometric lattice-lifting property which may be of independent interest. A Banach lattice X is said to have the *isometric lattice-lifting property* if there exists a lattice isometric embedding α from X to $\text{FBL}[X]$ such that $\widehat{id_X}\alpha = id_X$. This property was introduced in [3], inspired by previous works of Godefroy and Kalton [18] on Lipschitz-free spaces. Notably, Banach lattices ordered by a 1-unconditional basis have this property.

In [28, Theorem 8.3] an alternative proof of the lattice lifting property for spaces ordered by a 1-unconditional basis was given, which also worked with $\text{FBL}[E]$ replaced by $\text{FBL}^{(p)}[E]$. We now show how to generalize these results to $\text{FBL}^{\mathcal{C}}[E]$.

Proposition 3.7. *Let $X \in \overline{\mathcal{C}}$ be a Banach space with a normalized 1-unconditional basis (e_i) , viewed as a Banach lattice in the pointwise order induced by the basis. Then $\text{FBL}^{\mathcal{C}}[X]$ contains a lattice isometric copy of X . Moreover, there exists a contractive lattice homomorphic projection onto this sublattice.*

Proof. Using either [3, Theorem 4.1] or [28, Theorem 8.3] with $p = 1$ we obtain a lattice isometric embedding $\alpha : X \rightarrow \text{FBL}^{\mathcal{C}_1}[X]$. Since $\mathcal{C} \subseteq \mathcal{C}_1$, it follows that $\rho_{\mathcal{C}}(f) \leq \rho_{\mathcal{C}_1}(f)$ for every $f \in \text{FVL}[X]$, so the formal identity extends to a norm one lattice homomorphism $j : \text{FBL}^{\mathcal{C}_1}[X] \rightarrow \text{FBL}^{\mathcal{C}}[X]$. On the other hand, since $X \in \overline{\mathcal{C}}$, the identity operator on X extends to a contractive lattice homomorphism $\widehat{id_X} : \text{FBL}^{\mathcal{C}}[X] \rightarrow X$. It can be checked from the construction of α that $\widehat{id_X}j\alpha = id_X$, from which it follows that

$$\|x\|_X = \|\widehat{id_X}j\alpha x\| \leq \rho_{\mathcal{C}}(j\alpha x) \leq \rho_{\mathcal{C}_1}(\alpha x) \leq \|x\|_X$$

for every $x \in X$. □

Theorem 3.8. *The following are equivalent for a class of Banach lattices \mathcal{D} .*

- (1) *For any Banach space E , $\text{FBL}^{\mathcal{C}_p}[E] = \text{FBL}^{\mathcal{D}}[E]$ as sets and the norms $\rho_{\mathcal{C}_p}$ and $\rho_{\mathcal{D}}$ agree there.*
- (2) *$\mathcal{D} \subseteq \mathcal{C}_p$ and $\ell^p \in \overline{\mathcal{D}}$.*

Proof. (1) \implies (2): By Proposition 3.7, $\text{FBL}^{\mathcal{D}}[\ell^p] = \text{FBL}^{\mathcal{C}_p}[\ell^p]$ contains a lattice isometric copy of ℓ^p . By Proposition 2.7, $\text{FBL}^{\mathcal{D}}[\ell^p] \in \overline{\mathcal{D}}$. Hence, $\ell^p \in \overline{\mathcal{D}}$.

Suppose that $X \in \mathcal{D}$. Let $\phi_X : X \rightarrow \text{FBL}^{\mathcal{D}}[X]$ be the canonical embedding. The identity $i : X \rightarrow X$ induces a lattice homomorphic contraction $\widehat{i} : \text{FBL}^{\mathcal{D}}[X] \rightarrow X$ so that $\widehat{i}\phi_X = i$. Since $\text{FBL}^{\mathcal{D}}[X] = \text{FBL}^{\mathcal{C}_p}[X]$ lattice isometrically, this space belongs to $\overline{\mathcal{C}}_p = \mathcal{C}_p$ by Proposition 2.7. Hence,

so does $\text{FBL}^{\mathcal{D}}[X]/\ker \hat{i}$. Define $j : \text{FBL}^{\mathcal{D}}[X]/\ker \hat{i} \rightarrow X$ by $j[f] = \hat{i}f$, where $[f]$ is the equivalence class of f . The map j is a lattice isomorphism so that $\|j\| \leq 1$ and $x = j[\delta_x]$ for any $x \in X$. Hence, $B_X = j(B_{\text{FBL}^{\mathcal{D}}[X]/\ker \hat{i}})$. This shows that j is an onto lattice isometry. Therefore, $X \in \mathcal{C}_p$.

(2) \implies (1): Since $\mathcal{D} \subseteq \mathcal{C}_p$, for any Banach space E , $\rho_{\mathcal{D}} \leq \rho_{\mathcal{C}_p}$ on $\text{FVL}[E]$. Similarly, since $\ell^p \in \overline{\mathcal{D}}$, $\rho_{\{\ell^p\}} \leq \rho_{\overline{\mathcal{D}}}$. However, $\rho_{\mathcal{C}_p} = \rho_{\{\ell^p\}}$ and $\rho_{\overline{\mathcal{D}}} = \rho_{\mathcal{D}}$ by Corollary 2.6. Therefore, we have the reverse inequality $\rho_{\mathcal{C}_p} \leq \rho_{\mathcal{D}}$. \square

For the case of $\mathcal{C}_{p,\infty}$, the following result can be proven in essentially the same way as Theorem 3.8. The only difference is that, since $\ell^{p,\infty}$ does not have a basis, Proposition 3.7 cannot be applied directly. However, we can still use Proposition 3.7 to embed $\ell^{p,\infty}$ lattice isometrically into $(\bigoplus_m \text{FBL}^{\mathcal{C}_{p,\infty}}[\ell^{p,\infty}(m)])_{\infty}$ and continue with the rest of the proof with slight adaptations.

Theorem 3.9. *The following are equivalent for a class of Banach lattices \mathcal{D} .*

- (1) *There is a constant $M_1 < \infty$ so that for any Banach space E , $\text{FBL}^{\mathcal{C}_{p,\infty}}[E] = \text{FBL}^{\mathcal{D}}[E]$ as sets and $\frac{1}{M_1}\rho_{\mathcal{D}} \leq \rho_{\mathcal{C}_{p,\infty}} \leq M_1\rho_{\mathcal{D}}$ there.*
- (2) *There is a constant $M_2 < \infty$ so that any $X \in \mathcal{D}$ satisfies an upper p -estimate with constant M_2 , and $\ell^{p,\infty}$ is M_2 -lattice isomorphic to a space $Y \in \overline{\mathcal{D}}$.*

4. DISTINGUISHING $\rho_{\mathcal{X}_{p,\infty}}$ AND $\rho_{\mathcal{C}_{p,\infty}}$

A natural question that arises from Proposition 3.3 and Theorem 3.5 is whether the constant γ_p can be improved to 1. In this section, we will show that the answer is negative in general. In particular, there is a Banach space E so that $\rho_{\mathcal{C}_{p,\infty}} \neq \rho_{\mathcal{X}_{p,\infty}}$ on $\text{FVL}[E]$.

Let X, Y be Banach lattices and let $1 \leq C < \infty$. We say that X C -lattice embeds into Y if there exists a lattice isomorphic injection $T : X \rightarrow Y$ such that $C^{-1}\|x\| \leq \|Tx\| \leq \|x\|$ for all $x \in X$. In this case, we call T a C -lattice embedding. If $C = 1$, then T is a lattice isometric embedding. A Banach space with a normalized 1-unconditional basis is regarded as a Banach lattice in the coordinate-wise order.

Theorem 4.1. *Let X be a Banach lattice with a normalized 1-unconditional basis (e_i) . Denote the biorthogonal functionals by (e_i^*) . For any $C \geq 1$, the following statements are equivalent.*

- (1) *For each normalized finitely supported $a = \sum_{i=1}^n a_i e_i \in X_+$ and every $\varepsilon > 0$, there exist $b = \sum_{i=1}^n b_i e_i^* \in X_+$ and $d = (d_i)_{i=1}^n \in \mathbb{R}_+^n$ so that $\sum_{i=1}^n a_i b_i > 1 - \varepsilon$, $\sum_{i=1}^n d_i = 1$ and $\|\sum_{i \in I} b_i e_i^*\|^{p'} \leq C^{p'} \sum_{i \in I} d_i$ for any $I \subseteq \{1, \dots, n\}$.*
- (2) *There is a set Γ of (probability) measures such that X C -lattice embeds into $(\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu))_{\infty}$.*

Proof. (1) \implies (2): Let $a \in X_+$ be normalized and finitely supported, and let $\varepsilon > 0$ be given. Obtain b and d from (1). Let $(U_i)_{i=1}^n$ be a measurable partition of $(0, 1)$ such that $\lambda(U_i) = d_i$, $1 \leq i \leq n$, where λ denotes the Lebesgue measure. Define

$$S_{a,\varepsilon} : X \rightarrow L^{p,\infty}(0, 1) \text{ by } S_{a,\varepsilon}(\sum_{i=1}^{\infty} c_i e_i) = \sum_{i=1}^n \frac{c_i b_i \chi_{U_i}}{d_i}.$$

We have

$$\|S_{a,\varepsilon} a\| \geq \lambda(0, 1)^{-\frac{1}{p'}} \int_0^1 |S_{a,\varepsilon} a| d\lambda = \sum_{i=1}^n a_i b_i > 1 - \varepsilon.$$

On the other hand, if $\|\sum_{i=1}^{\infty} c_i e_i\| \leq 1$ then there exists $I \subseteq \{1, \dots, n\}$ so that

$$\begin{aligned} \|S_{a,\varepsilon}c\| &= \lambda\left(\bigcup_{i \in I} U_i\right)^{-\frac{1}{p'}} \int_{\bigcup_{i \in I} U_i} |S_{a,\varepsilon}c| d\lambda = \left(\sum_{i \in I} d_i\right)^{-\frac{1}{p'}} \sum_{i \in I} |c_i| b_i \\ &= \left(\sum_{i \in I} d_i\right)^{-\frac{1}{p'}} \left\langle \sum_{i=1}^{\infty} |c_i| e_i, \sum_{i \in I} b_i e_i^* \right\rangle \leq \left(\sum_{i \in I} d_i\right)^{-\frac{1}{p'}} \left\| \sum_{i \in I} b_i e_i^* \right\| \leq C. \end{aligned}$$

Let Γ be the Cartesian product of the set of normalized finitely supported elements in X_+ and the interval $(0, \frac{1}{2})$. It is now clear that the map $T : X \rightarrow (\bigoplus_{(a,\varepsilon) \in \Gamma} L^{p,\infty}(0,1))_{\infty}$ given by

$$T\left(\sum_{i=1}^{\infty} c_i e_i\right) = (C^{-1}S_{a,\varepsilon}c)_{(a,\varepsilon) \in \Gamma}$$

is a C -lattice embedding.

(2) \implies (1): Let $T : X \rightarrow (\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu))_{\infty}$ be a C -lattice embedding. Suppose that $a = \sum_{i=1}^n a_i e_i$ is normalized and $\varepsilon > 0$. For each $\mu_0 \in \Gamma$, let P_{μ_0} be the projection of $(\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu))_{\infty}$ onto the μ_0 -th component. There exists μ_0 such that $C\|P_{\mu_0}Ta\| > 1 - \varepsilon$. Let $f_i = P_{\mu_0}T e_i$. Choose a μ_0 -measurable set U such that $0 < \mu_0(U) < \infty$ and

$$C \int_U P_{\mu_0}Ta d\mu_0 > (1 - \varepsilon)(\mu_0(U))^{\frac{1}{p'}}.$$

Set $U_i = U \cap \text{supp } f_i$ and $m = (\sum_{i=1}^n \mu_0(U_i))^{\frac{1}{p'}}$. Define

$$b_i = \frac{C \int_{U_i} f_i d\mu_0}{m}, \quad d_i = \frac{\mu_0(U_i)}{m^{p'}}, \quad 1 \leq i \leq n.$$

We have $d_i \geq 0$, $\sum_{i=1}^n d_i = 1$ and

$$\sum_{i=1}^n a_i b_i = \frac{C}{m} \int_{\bigcup_{i=1}^n U_i} \sum_{i=1}^n a_i f_i d\mu_0 = \frac{C}{m} \int_{\bigcup_{i=1}^n U_i} P_{\mu_0}Ta > 1 - \varepsilon.$$

On the other hand, suppose that $I \subseteq \{1, \dots, n\}$. If $\|\sum_{i=1}^{\infty} c_i e_i\| \leq 1$, then

$$\begin{aligned} \left\langle \sum_{i=1}^{\infty} c_i e_i, \sum_{i \in I} b_i e_i^* \right\rangle &= \sum_{i \in I} c_i b_i = \frac{C}{m} \int_{\bigcup_{i \in I} U_i} \sum_{i \in I} c_i f_i d\mu_0 \\ &\leq \frac{C}{m} \|P_{\mu_0}T(\sum_{i=1}^{\infty} c_i e_i)\| (\mu_0(\bigcup_{i \in I} U_i))^{\frac{1}{p'}} \leq C \left(\sum_{i \in I} d_i\right)^{\frac{1}{p'}}. \end{aligned}$$

This proves that $\|\sum_{i \in I} b_i e_i^*\|^{p'} \leq C^{p'} \sum_{i \in I} d_i$. \square

Remark 4.2. If X is a two dimensional Banach lattice that satisfies an upper p -estimate with constant one, then $C = 1$ works in Theorem 4.1. Indeed, let (e_1, e_2) be a normalized 1-unconditional basis for X . Suppose that $a = a_1 e_1 + a_2 e_2$ is normalized and non-negative. Choose $b = b_1 e_1^* + b_2 e_2^* \in X_+^*$ such that $\langle a, b \rangle = 1 = \|b\|$. Define $d_i = b_i^{p'} / (b_1^{p'} + b_2^{p'})$, $i = 1, 2$. Note that X^* satisfies a lower p' -estimate with constant one and e_1^* and e_2^* are disjoint, so

$$b_1^{p'} + b_2^{p'} = \|b_1 e_1^*\|^{p'} + \|b_2 e_2^*\|^{p'} \leq \|b\|^{p'} = 1.$$

It is easy to check that $\|\sum_{i \in I} b_i e_i^*\|^{p'} \leq \sum_{i \in I} d_i$ for any $I \subseteq \{1, 2\}$ and $d_1 + d_2 = 1$.

Corollary 4.3. *Suppose that X is a Banach lattice with a normalized 1-unconditional basis (e_i) that C -lattice embeds into $(\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu))_\infty$. For any normalized $b = \sum_{i=1}^n b_i e_i^* \in X_+^*$ and any $I_j \subseteq \{1, \dots, n\}$, $1 \leq j \leq m$, such that $\sum_{j=1}^m \chi_{I_j} = l \chi_{\{1, \dots, n\}}$ for some $l \in \mathbb{N}$, we have*

$$\sum_{j=1}^m \left\| \sum_{i \in I_j} b_i e_i^* \right\|^{p'} \leq C^{p'} l.$$

Proof. Choose a normalized $a = \sum_{i=1}^n a_i e_i \in X_+$ so that $\sum_{i=1}^n a_i b_i = 1$. First assume that $[(e_i)_{i=1}^n]$ is uniformly convex. Let $\varepsilon > 0$ be given. There exists $\delta > 0$ so that if $b' = \sum_{i=1}^n b'_i e_i^* \in X_+^*$, $\|b'\| = 1$ and $\langle a, b' \rangle > 1 - \delta$, then $\|b - b'\| < \varepsilon$. By Theorem 4.1, there exists $b' \in X_+^*$ and $d = (d_i)_{i=1}^n \in \mathbb{R}_+^n$ so that $\langle a, b' \rangle > 1 - \delta$ and $\|\sum_{i \in I} b'_i e_i^*\|^{p'} \leq C^{p'} \sum_{i \in I} d_i$ for any $I \subseteq \{1, \dots, n\}$. It follows that for any $I \subseteq \{1, \dots, n\}$,

$$\left\| \sum_{i \in I} b_i e_i^* \right\| \leq \left\| \sum_{i \in I} |b_i - b'_i| e_i^* \right\| + \left\| \sum_{i \in I} b'_i e_i^* \right\| \leq \varepsilon \cdot \#I + C \left(\sum_{i \in I} d_i \right)^{\frac{1}{p'}}.$$

Now assume that $I_j \subseteq \{1, \dots, n\}$, $1 \leq j \leq m$, and that $\sum_{j=1}^m \chi_{I_j} = l \chi_{\{1, \dots, n\}}$ for some $l \in \mathbb{N}$. Set $A_j = \varepsilon \cdot \#I_j$ and $B_j = C \left(\sum_{i \in I_j} d_i \right)^{\frac{1}{p'}}$. Then

$$\begin{aligned} \left(\sum_{j=1}^m \left\| \sum_{i \in I_j} b_i e_i^* \right\|^{p'} \right)^{\frac{1}{p'}} &\leq \left(\sum_{j=1}^m (A_j + B_j)^{p'} \right)^{\frac{1}{p'}} \leq \left(\sum_{j=1}^m A_j^{p'} \right)^{\frac{1}{p'}} + \left(\sum_{j=1}^m B_j^{p'} \right)^{\frac{1}{p'}} \\ &\leq \sum_{j=1}^m A_j + \left(\sum_{j=1}^m B_j^{p'} \right)^{\frac{1}{p'}} = \varepsilon \sum_{j=1}^m \#I_j + C \left(\sum_{j=1}^m \sum_{i \in I_j} d_i \right)^{\frac{1}{p'}} \leq \varepsilon n l + C l^{\frac{1}{p'}}. \end{aligned}$$

Taking $\varepsilon \downarrow 0$ gives the desired result.

We now return to a general X with a normalized 1-unconditional basis (e_i) . Given a normalized $b = \sum_{i=1}^n b_i e_i^* \in X_+^*$ and $\varepsilon > 0$, there is a uniformly convex lattice norm $||| \cdot |||$ on $[(e_i)_{i=1}^n]$ such that $\|x\| \leq |||x||| \leq (1 + \varepsilon)\|x\|$ for any $x \in [(e_i)_{i=1}^n]$. Set $u_i = e_i / |||e_i|||$, $u_i^* = e_i^* / |||e_i^*|||$ and $b' = \sum_{i=1}^n b'_i u_i^* = b / |||b|||$. Note that $(u_i)_{i=1}^n$ is a 1-unconditional basis for $[(e_i)_{i=1}^n] = [(u_i)_{i=1}^n]$ and b' is $||| \cdot |||$ -normalized. Moreover, if $T : [(e_i)_{i=1}^n] \rightarrow (\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu))_\infty$ is a C -lattice embedding, then

$$\tilde{T}x = Tx : [(u_i)_{i=1}^n] \rightarrow \left(\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\mu) \right)_\infty$$

is a $(1 + \varepsilon)C$ -lattice embedding. Assume that $I_j \subseteq \{1, \dots, n\}$, $1 \leq j \leq m$, such that $\sum_{j=1}^m \chi_{I_j} = l \chi_{\{1, \dots, n\}}$ for some $l \in \mathbb{N}$. By the first part,

$$\sum_{j=1}^m \left\| \sum_{i \in I_j} b'_i u_i^* \right\|^{p'} \leq (1 + \varepsilon)^{p'} C^{p'} l.$$

Since $|||b||| \leq 1 + \varepsilon$, for any $I \subseteq \{1, \dots, n\}$ we have

$$\left\| \sum_{i \in I} b'_i u_i^* \right\| = \frac{1}{|||b|||} \left\| \sum_{i \in I} b_i e_i^* \right\| \geq \frac{1}{1 + \varepsilon} \left\| \sum_{i \in I} b_i e_i^* \right\|.$$

Thus,

$$\sum_{j=1}^m \left\| \sum_{i \in I_j} b_i e_i^* \right\|^{p'} \leq (1 + \varepsilon)^{2p'} C^{p'} l.$$

Taking $\varepsilon \downarrow 0$ gives the desired result. \square

The next example shows that γ_p cannot be made to be 1 in Proposition 3.3.

Example 4.4. *Suppose that $1 < p < \infty$. Denote by C_p the infimum of the set of constants C so that every (p, ∞) -convex Banach lattice with constant one C -lattice embeds into $(\bigoplus_{\gamma \in \Gamma} L^{p, \infty}(\mu))_\infty$ for some set of measures Γ . Then $C_p^{p'} \geq \frac{3 \cdot 2^{p'}}{2(1+2^{p'})} > 1$.*

Proof. Let $X = \mathbb{R}^3$ so that the norm on X^* is

$$\|(b_1, b_2, b_3)\|_{X^*} = \max\left\{\left(|b_i|^{p'} + (|b_j| + |b_k|)^{p'}\right)^{\frac{1}{p'}} : \{i, j, k\} = \{1, 2, 3\}\right\}.$$

It is clear that the coordinate unit vectors form a normalized 1-unconditional basis for X . Suppose that $b = (b_1, b_2, b_3) \in X^*$ and I, J are disjoint non-empty subsets of $\{1, 2, 3\}$. Without loss of generality, we may assume that $I = \{i, j\}$ and $J = \{k\}$, where i, j, k are distinct. Then

$$\|b\chi_I\|_{X^*}^{p'} + \|b\chi_J\|_{X^*}^{p'} = (|b_i| + |b_j|)^{p'} + |b_k|^{p'} \leq \|b\|_{X^*}^{p'}.$$

Thus, X^* satisfies a lower p' -estimate with constant 1. Hence, X satisfies an upper p -estimate with constant 1. Suppose that X C -lattice embeds into some $(\bigoplus_{\gamma \in \Gamma} L^{p, \infty}(\mu))_\infty$. The vector $(1 + 2^{p'})^{-\frac{1}{p'}}(1, 1, 1)$ is normalized in X_+^* . By Corollary 4.3,

$$\frac{1}{1 + 2^{p'}} (\|(1, 1, 0)\|^{p'} + \|(1, 0, 1)\|^{p'} + \|(0, 1, 1)\|^{p'}) \leq 2C^{p'}.$$

Thus,

$$\frac{3 \cdot 2^{p'}}{2(1 + 2^{p'})} \leq C^{p'}.$$

Taking infimum over the eligible C 's gives the desired result. Direct verification shows that $\frac{3 \cdot 2^{p'}}{2(1+2^{p'})} > 1$ if $1 < p < \infty$. \square

Now we are ready to show that γ_p also cannot be made to be 1 in Theorem 3.5. Note that the following result can be adapted to work for other classes of lattices \mathcal{C} .

Proposition 4.5. *Let X be a Banach lattice with a normalized 1-unconditional basis (e_i) satisfying upper p -estimates. Let $C \geq 1$. The following are equivalent.*

- (1) X C -lattice embeds in $(\bigoplus_{\gamma \in \Gamma} L^{p, \infty}(\mu))_\infty$.
- (2) For every operator $T : X \rightarrow X$, the extension $\widehat{T} : \text{FBL}^{\mathcal{X}_{p, \infty}}[X] \rightarrow X$ satisfies $\|\widehat{T}\| \leq C\|T\|$.
- (3) The extension $\widehat{id}_X : \text{FBL}^{\mathcal{X}_{p, \infty}}[X] \rightarrow X$ of the identity $id_X : X \rightarrow X$ satisfies $\|\widehat{id}_X\| \leq C$.

Proof. Clearly, (1) \Rightarrow (2) \Rightarrow (3).

For the proof of (3) \Rightarrow (1) we first note that by the remark after Theorem 3.5, for any Banach space E , $\rho_{\mathcal{X}_{p, \infty}^\sigma} = \rho_{\{\ell^{p, \infty}\}}$ on $\text{FVL}[E]$. Thus, it follows from Proposition 2.7 that $\text{FBL}^{\mathcal{X}_{p, \infty}}[E]$ belongs to the class $\{\widehat{\ell^{p, \infty}}\}$, i.e., it lattice isometrically embeds into $(\bigoplus_{\gamma \in \Gamma} \ell^{p, \infty})_\infty$.

We now adapt the proof of Proposition 3.7 to our setting. Recall that by [28, Theorem 8.3] there is a lattice isometric embedding $\alpha : X \rightarrow \text{FBL}[X] = \text{FBL}^{C_1}[X]$ such that $\widehat{id}_X \alpha = id_X$. Moreover, since $\mathcal{X}_{p, \infty}^\sigma \subseteq \mathcal{C}_1$, the formal identity $j : \text{FBL}^{C_1}[X] \rightarrow \text{FBL}^{\mathcal{X}_{p, \infty}^\sigma}[X]$ is a continuous norm one injection such that $\widehat{id}_X j \alpha = id_X$. We claim that $j \alpha : X \rightarrow \text{FBL}^{\mathcal{X}_{p, \infty}^\sigma}[X]$ defines a C -lattice embedding. If

true, the previous paragraph gives the implication (3) \Rightarrow (1). To prove the claim, simply note that for every $x \in X$

$$\frac{1}{C}\|x\| \leq \frac{1}{\|\widehat{id_X}\|} \|\widehat{id_X} j\alpha x\| \leq \rho_{\mathcal{X}_{p,\infty}}(j\alpha x) \leq \rho_{\mathcal{C}_1}(\alpha x) \leq \|x\|.$$

□

4.1. Equivalent renormings of $L^{p,\infty}(\mu)$. Let $1 < p < \infty$. As mentioned above, given a measure space (Ω, Σ, μ) , the space $L^{p,\infty}(\mu)$ is the set of all measurable functions $f : \Omega \rightarrow \mathbb{R}$ such that

$$\|f\|_{L^{p,\infty}}^* := \sup_{t>0} t\mu(\{|f| > t\})^{\frac{1}{p}}$$

is finite. In the previous subsections, we equipped $L^{p,\infty}(\mu)$ with the norm

$$\|f\|_{L^{p,\infty}} = \sup\{\mu(E)^{\frac{1}{p}-1} \int_E |f| d\mu : 0 < \mu(E) < \infty\}.$$

However, for each $1 \leq r < p$, we could have equally chosen to equip $L^{p,\infty}(\mu)$ with the norm

$$\|f\|_{L_r^{p,\infty}} := \sup_{0 < \mu(A) < \infty} \mu(A)^{-\frac{1}{r} + \frac{1}{p}} \left(\int_A |f|^r d\mu \right)^{\frac{1}{r}}.$$

Indeed, it is well-known (cf. [19, Exercise 1.1.12]) that

$$\|f\|_{L^{p,\infty}}^* \leq \|f\|_{L_r^{p,\infty}} \leq \left(\frac{p}{p-r} \right)^{\frac{1}{r}} \|f\|_{L^{p,\infty}}^*$$

for every $f \in L^{p,\infty}(\mu)$, and it is easy to see that for each $1 \leq r < p$, $\|\cdot\|_{L_r^{p,\infty}}$ defines a lattice norm on $L^{p,\infty}(\mu)$ with upper p -estimate constant 1. For this reason, it is natural to ask whether the ambiguity in choosing a norm on $L^{p,\infty}(\mu)$ is the reason why γ_p cannot be chosen to be 1 in Proposition 3.3 and Theorem 3.5.

The aim of this subsection is to show that $(L^{p,\infty}(\mu), \|\cdot\|_{L_r^{p,\infty}})$ lattice isometrically embeds into a Banach lattice of the form $(\bigoplus_{\mu \in \Gamma} L^{p,\infty}(\nu))_\infty$, where each $L^{p,\infty}(\nu)$ is given the $L_1^{p,\infty}$ -norm. As a consequence, we deduce that $\rho_{p,\infty}(\cdot) = \rho_{L_1^{p,\infty}}(\cdot) \geq \rho_{L_r^{p,\infty}}(\cdot)$ on $\text{FVL}[E]$, proving that the above renormings cannot eliminate the constant γ_p in Proposition 3.3 and Theorem 3.5.

We begin with a lemma.

Lemma 4.6. *Let $e = (1, \dots, 1) \in \mathbb{R}^n$ and let $(e_i)_{i=1}^n$ be the coordinate unit vector basis for \mathbb{R}^n . Let $\beta, b \in \mathbb{R}_+^n$, $\|\beta\|_1, \|b\|_1 \leq 1$, and let $0 < s < 1$. Define $\alpha : \mathbb{R}_+^n \rightarrow \mathbb{R}$ by $\alpha(x) = (\beta \cdot x)^{1-s} (b \cdot x)^s$, where \cdot is the dot product on \mathbb{R}^n . Then α is positively homogeneous, $\alpha(e) = \|\beta\|_1^{1-s} \cdot \|b\|_1^s \leq 1$ and $\sum_{j=1}^m \alpha(x_j) \leq \alpha(\sum_{j=1}^m x_j)$ for any sequence $(x_j)_{j=1}^m \subseteq \ell^1(n)_+$.*

Proof. Let $(x_j)_{j=1}^m \subseteq \mathbb{R}_+^n$. Then

$$\begin{aligned} \sum_{j=1}^m \alpha(x_j) &\leq \|(\beta \cdot x_j)^{1-s}\|_{\frac{1}{1-s}} \| (b \cdot x_j)^s \|_{\frac{1}{s}} \\ &= (\beta \cdot \sum_{j=1}^m x_j)^{1-s} (b \cdot \sum_{j=1}^m x_j)^s = \alpha(\sum_{j=1}^m x_j). \end{aligned}$$

The remaining assertions are clear. □

Let $x \in \mathbb{R}^n$ be expressed as $(x(j))_{j=1}^n$. In our next result, for fixed $i \in \mathbb{N}$, we write $x < i$ if $\max \text{supp } x < i$.

Proposition 4.7. *There is a sequence $(d_i)_{i=1}^n$ of non-negative real numbers such that for each i ,*

$$\sum_{j=1}^{i-1} x(j)d_j + \alpha(z) - m - \sum_{j=1}^{i-1} y(j)d_j \leq d_i \leq m' + \sum_{j=1}^{i-1} y'(j)d_j - \sum_{j=1}^{i-1} x'(j)d_j - \alpha(z')$$

provided that $m, m' \in \mathbb{R}_+$, $x, y, z, x', y', z' \in \mathbb{R}_+^n$,

$$(4.1) \quad x, y, x', y' < i, \quad x + z = e_i + me + y \quad \text{and} \quad e_i + x' + z' = m'e + y'.$$

Proof. The proof is by induction on i . For $i = 1$, $x = y = x' = y' = 0$. Thus it suffices to show that

$$\alpha(z) - m \leq m' - \alpha(z') \quad \text{if} \quad z = e_1 + me \quad \text{and} \quad e_1 + z' = m'e.$$

To see this, note that

$$m + m' \geq \alpha((m + m')e) = \alpha(z + z') \geq \alpha(z) + \alpha(z'),$$

from which the desired conclusion follows.

Assume now that d_j , $1 \leq j < i$, have been chosen to satisfy the proposition. To be able to choose d_i satisfying the proposition, we need to show that

$$\sum_{j=1}^{i-1} x(j)d_j + \alpha(z) - m - \sum_{j=1}^{i-1} y(j)d_j \leq m' + \sum_{j=1}^{i-1} y'(j)d_j - \sum_{j=1}^{i-1} x'(j)d_j - \alpha(z')$$

if $m, m', x, y, z, x', y', z'$ satisfy (4.1). Equivalently, we must show that

$$(4.2) \quad \sum_{j=1}^{i-1} (x(j) + x'(j))d_j + \alpha(z) + \alpha(z') \leq m + m' + \sum_{j=1}^{i-1} (y(j) + y'(j))d_j.$$

If $x + x' = y + y'$, then (4.1) gives $z + z' = (m + m')e$. Thus, $m + m' \geq \alpha(z) + \alpha(z')$, yielding (4.2). Otherwise, let $u = (x + x') \wedge (y + y') \in \mathbb{R}_+^n$. Then $u < i$ and $v_1 := x + x' - u$, $v_2 := y + y' - u$ are disjoint in \mathbb{R}_+^n , not both 0. Assume that $j_0 = \max \text{supp } v_1 > \max \text{supp } v_2$, as the other case is similar. Set $w = v_1 - v_1(j_0)e_{j_0}$. Then $w, v_2 < j_0$, $w(j) = v_1(j)$ if $j < j_0$ and

$$v_1(j_0)e_{j_0} + w + (z + z') = (m + m')e + v_2.$$

Using the right inequality in the inductive hypothesis and the positive homogeneity of α , we see that

$$v_1(j_0)d_{j_0} \leq (m + m') + \sum_{j=1}^{j_0-1} v_2(j)d_j - \sum_{j=1}^{j_0-1} v_1(j)d_j - \alpha(z + z').$$

Adding $\sum_{j=1}^{j_0-1} u(j)d_j$ to both sides and rearranging gives

$$\sum_{j=1}^{i-1} (x(j) + x'(j))d_j + \alpha(z + z') \leq m + m' + \sum_{j=1}^{i-1} (y(j) + y'(j))d_j.$$

The desired inequality (4.2) then follows since $\alpha(z) + \alpha(z') \leq \alpha(z + z')$. \square

Corollary 4.8. *In the above notation, we have $d = (d_i)_{i=1}^n \in \mathbb{R}_+^n$, $\alpha(e) \leq \|d\|_1 \leq 1$ and $\alpha(x) \leq x \cdot d$ for any $x \in \mathbb{R}_+^n$.*

Proof. Taking $i = n$ in Proposition 4.7, we have

$$x = 0, y = e - e_n, z = e, m = 0; x' = e - e_n, y' = 0, z' = 0, m' = 1.$$

Note that $x, y, x', y' < n$, $x + z = e_n + me + y$ and $e_n + x' + z' = m'e + y'$. Thus,

$$\alpha(e) - \sum_{j=1}^{n-1} d_j \leq d_n \leq 1 - \sum_{j=1}^{n-1} d_j.$$

Hence, $\alpha(e) \leq \|d\|_1 \leq 1$. Finally, if $u \in \mathbb{R}_+^n$, let $x = 0$, $z = u$, $m = 0$ and $y = u - u(n)e_n$. Then $x + z = u(n)e_n + me + y$. By Proposition 4.7 and the positive homogeneity of α ,

$$\alpha(u) - \sum_{j=1}^{n-1} u(j)d_j \leq u(n)d_n \implies \alpha(u) \leq u \cdot d.$$

□

Lemma 4.9. *Let (Ω, Σ, μ) be a measure space. Suppose that $(a_i)_{i=1}^n \in \mathbb{R}_+^n$ and $(U_i)_{i=1}^n$ is a disjoint sequence in Σ with $0 < \mu(U_i) < \infty$ so that $C := (\sum_{i=1}^n \mu(U_i))^{-\frac{1}{r} + \frac{1}{p}} \|\sum_{i=1}^n a_i \chi_{U_i}\|_{L^r(\mu)} \leq 1$. There is a measure ν on Σ and a linear lattice homomorphism $S : L_r^{p,\infty}(\mu) \rightarrow L_1^{p,\infty}(\nu)$ so that $\|S\| \leq 1$ and $\|S(\sum_{i=1}^n a_i \chi_{U_i})\| \geq C^r$.*

Proof. Define $M = \sum_{j=1}^n \mu(U_j)$, $b_i = \frac{\mu(U_i)}{M}$, $b = (b_i)_{i=1}^n$ and $\beta_i = M^{\frac{r}{p}-1} \mu(U_i) a_i^r$, $\beta = (\beta_i)_{i=1}^n$. Clearly, $\beta, b \in \mathbb{R}_+^n$ and $\|b\|_1 = 1$. Moreover,

$$\|\beta\|_1 = M^{r(\frac{1}{p}-\frac{1}{r})} \|\sum_{i=1}^n a_i \chi_{U_i}\|_{L^r(\mu)}^r = C^r \leq 1.$$

Take $s = p'(\frac{1}{r} - \frac{1}{p})$ and define α as in Lemma 4.6. Note that $\alpha(e) = \|\beta\|_1^{1-s} \|b\|_1^s = C^{r(1-s)} \leq 1$. Obtain d using Proposition 4.7. Define a measure ν on Σ by $\nu(A) = \sum_{i=1}^n \frac{d_i \mu(A \cap U_i)}{\mu(U_i)}$ and a linear operator

$$S : L_r^{p,\infty}(\mu) \rightarrow L_1^{p,\infty}(\nu) \quad \text{by} \quad Sf = M^{\frac{r}{p}} \sum_{i=1}^n \frac{a_i^{r-1} b_i}{d_i} f \chi_{U_i}.$$

Clearly, S is a linear lattice homomorphism. Note that $\nu(\cup_{i=1}^n U_i) = \sum_{i=1}^n d_i \leq 1$. Hence,

$$\begin{aligned} \|S(\sum_{i=1}^n a_i \chi_{U_i})\| &\geq M^{\frac{r}{p}} \|\sum_{i=1}^n \frac{a_i^r b_i}{d_i} \chi_{U_i}\|_{L^1(\nu)} = M^{\frac{r}{p}} \sum_{i=1}^n a_i^r b_i \\ &= M^{\frac{r}{p}-1} \|\sum_{i=1}^n a_i \chi_{U_i}\|_{L^r(\mu)}^r = C^r. \end{aligned}$$

On the other hand, let $f \in L_r^{p,\infty}(\mu)$ with $\|f\| \leq 1$. If $A \in \Sigma$ with $0 < \nu(A) < \infty$ then

$$\begin{aligned} \int_A |Sf| d\nu &= M^{\frac{r}{p}} \sum_{i=1}^n \frac{a_i^{r-1} b_i}{d_i} \int_{A \cap U_i} |f| d\nu = M^{\frac{r}{p}-1} \sum_{i=1}^n a_i^{r-1} \int_{A \cap U_i} |f| d\mu \\ &\leq M^{\frac{r}{p}-1} \left\| \sum_{i=1}^n a_i^{r-1} \chi_{A \cap U_i} \right\|_{L^{r'}(\mu)} \left\| f \chi_{\cup_{i=1}^n (A \cap U_i)} \right\|_{L^r(\mu)} \\ &\leq \left(\sum_{i=1}^n \beta_i \frac{\mu(A \cap U_i)}{\mu(U_i)} \right)^{\frac{1}{r'}} \left(\sum_{i=1}^n b_i \frac{\mu(A \cap U_i)}{\mu(U_i)} \right)^{\frac{1}{r} - \frac{1}{p}}. \end{aligned}$$

Let $x = \left(\frac{\mu(A \cap U_i)}{\mu(U_i)}\right)_{i=1}^n \in \mathbb{R}_+^n$. By direct calculation, $1 - s = \frac{p'}{r'}$. Thus,

$$\left| \int_A S f d\nu \right| \leq (\beta \cdot x)^{\frac{1-s}{p'}} (b \cdot x)^{\frac{s}{p'}} = \alpha(x)^{\frac{1}{p'}} \leq (x \cdot d)^{\frac{1}{p'}} = \nu(A)^{\frac{1}{p'}}.$$

Hence, $\|Sf\| \leq 1$. □

Theorem 4.10. *Suppose that (Ω, Σ, μ) is a measure space and $1 < r < p$. There is a set Γ of measures on Σ so that $L_r^{p,\infty}(\mu)$ lattice isometrically embeds into $(\bigoplus_{\nu \in \Gamma} L_1^{p,\infty}(\nu))_\infty$.*

Proof. Let Γ be the set of simple functions $f = \sum_{i=1}^n a_i \chi_{U_i}$, where $n \in \mathbb{N}$, $a_i > 0$, $0 < \mu(U_i) < \infty$ and $(U_i)_{i=1}^n$ is a disjoint sequence in Σ with

$$C_f = \left(\sum_{i=1}^n \mu(U_i) \right)^{-\frac{1}{r} + \frac{1}{p}} \left\| \sum_{i=1}^n a_i \chi_{U_i} \right\|_r \leq 1.$$

For each $f \in \Gamma$, use the above results to choose a lattice homomorphism $S : L_r^{p,\infty}(\mu) \rightarrow L_1^{p,\infty}(\nu_f)$ so that $\|S_f\| \leq 1$ and $\|S_f f\| \geq C_f^r$. Define

$$T : L_r^{p,\infty}(\mu) \rightarrow \left(\bigoplus_{f \in \Gamma} L_1^{p,\infty}(\nu_f) \right)_\infty \quad \text{by} \quad Tg = (S_f g)_{f \in \Gamma}.$$

Clearly, T is a lattice homomorphism and $\|T\| \leq 1$. Suppose that $g \in L_r^{p,\infty}(\mu)_+$ with $\|g\|_{L_r^{p,\infty}} = 1$. For any $\varepsilon > 0$, there exists a non-negative simple function h such that $g \geq h$ and $\|h\|_{L_r^{p,\infty}} \geq 1 - \varepsilon$. Write $h = \sum_{i=1}^n a_i \chi_{U_i}$. There exists $I \subseteq \{1, \dots, n\}$ so that

$$\left\| \left(\sum_{i \in I} \mu(U_i) \right)^{-\frac{1}{r} + \frac{1}{p}} \sum_{i \in I} a_i \chi_{U_i} \right\|_r = \|h\|_{L_r^{p,\infty}} \geq 1 - \varepsilon.$$

Then $f := \sum_{i \in I} a_i \chi_{U_i} \in \Gamma$ with $C_f \geq 1 - \varepsilon$. Thus,

$$\|Tg\| \geq \|S_f g\| \geq \|S_f f\| \geq C_f^r \geq (1 - \varepsilon)^r,$$

so that $\|Tg\| \geq 1$. This shows that T is an isometric embedding. □

5. SUBSPACE PROBLEM

In this section we characterize when an embedding $\iota : F \hookrightarrow E$ induces a lattice embedding $\bar{\iota} : \text{FBL}^{(p,\infty)}[F] \hookrightarrow \text{FBL}^{(p,\infty)}[E]$. The analogous problem for $\text{FBL}^{(p)}$ was solved in [28, Theorem 3.7] by making use of an extension theorem for regular operators due to Pisier [31, Theorem 4]. This scheme of proof, however, does not seem to be applicable to $\text{FBL}^{(p,\infty)}$. For this reason, we develop in Section 5.1 an entirely new approach to the subspace problem which is based on push-outs. Then, in Section 5.2 we prove the injectivity of ℓ^p in the class of p -convex Banach lattices, which shows that our solution to the subspace problem for $\text{FBL}^{(p)}$ is equivalent to the one in [28]. Finally, in Section 5.3 we prove that $\ell^{p,\infty}$ is *not* injective in the class of Banach lattices with upper p -estimates.

We begin with some preliminaries. Let F be a closed subspace of a Banach space E and let $\iota : F \rightarrow E$ be the inclusion map. Let $q = \iota^* : E^* \rightarrow F^*$ and note that q is weak*-to-weak* continuous with $q(B_{E^*}) = B_{F^*}$. Hence, $f \circ q \in C(B_{E^*})$ if $f \in \text{FVL}[F]$.

Proposition 5.1. *The map $\bar{\iota} : \text{FVL}[F] \rightarrow \text{FVL}[E]$, $\bar{\iota}f = f \circ q$, is a vector lattice isomorphism from $\text{FVL}[F]$ onto the sublattice \mathcal{L} of $\text{FVL}[E]$ generated by $\{\delta_{\iota y} : y \in F\}$.*

Proof. Clearly, $\bar{\tau}$ is a lattice homomorphism from $\text{FVL}[F]$ into $C(B_{E^*})$. Moreover, $(\bar{\tau}\delta_y)(x^*) = (\delta_y \circ q)(x^*) = \delta_{\iota y}(x^*)$ for all $y \in F$ and $x^* \in B_{E^*}$. Hence, $\bar{\tau}\delta_y = \delta_{\iota y}$. Thus, $\bar{\tau}\text{FVL}[F]$ is a sublattice of $C(B_{E^*})$ that contains $\delta_{\iota y}$ for all $y \in F$, which implies that $\mathcal{L} \subseteq \bar{\tau}\text{FVL}[F]$. On the other hand, $\bar{\tau}^{-1}\mathcal{L}$ is a sublattice of $\text{FVL}[F]$ containing δ_y for all $y \in F$. Hence, $\bar{\tau}^{-1}\mathcal{L} = \text{FVL}[F]$, which implies that $\bar{\tau}\text{FVL}[F] = \mathcal{L}$. \square

Note that $\phi_{E\iota} : F \rightarrow \text{FBL}^{\mathcal{C}}[E]$ is a contraction (in fact, an isometry) and $\text{FBL}^{\mathcal{C}}[E] \in \bar{\mathcal{C}}$. By Proposition 2.5, $\widehat{\phi_{E\iota}} : (\text{FVL}[F], \rho_{\mathcal{C}}) \rightarrow \text{FBL}^{\mathcal{C}}[E]$ is a contraction.

Proposition 5.2. *Let $\bar{\tau}$ be the map from Proposition 5.1. Then $\widehat{\phi_{E\iota}}f = \bar{\tau}f$ for any $f \in \text{FVL}[F]$.*

Proof. Let $f \in \text{FVL}[F]$. Suppose that $f = G(\delta_{y_1}, \dots, \delta_{y_n})$, where G is a lattice-linear expression and $y_1, \dots, y_n \in F$. For any $x^* \in B_{E^*}$,

$$\begin{aligned} (\bar{\tau}f)(x^*) &= f(qx^*) = G(qx^*(y_1), \dots, qx^*(y_n)) = G(x^*(\iota y_1), \dots, x^*(\iota y_n)) \\ &= G(\delta_{\iota y_1}, \dots, \delta_{\iota y_n})(x^*). \end{aligned}$$

Thus, $\bar{\tau}f = G(\delta_{\iota y_1}, \dots, \delta_{\iota y_n})$. On the other hand,

$$\widehat{\phi_{E\iota}}(f) = G(\phi_{E\iota}y_1, \dots, \phi_{E\iota}y_n) = G(\delta_{\iota y_1}, \dots, \delta_{\iota y_n}).$$

Therefore, $\bar{\tau} = \widehat{\phi_{E\iota}}$. \square

The operator $\bar{\tau}$ extends to a contraction from $\text{FBL}^{\mathcal{C}}[F]$ into $\text{FBL}^{\mathcal{C}}[E]$, which we denote again by $\bar{\tau}$. The objective now is to determine when $\bar{\tau}$ is an embedding. We begin with the following elementary observation.

Proposition 5.3. *Suppose that for any $X \in \mathcal{C}$, any contraction $T : F \rightarrow X$ extends to a contraction $S : E \rightarrow X$. Then $\bar{\tau}$ is a lattice isometric embedding.*

Proof. In view of the preceding propositions and the fact that $\widehat{\phi_{E\iota}}$ is a contraction, it suffices to show that $\rho_{\mathcal{C}}(f) \leq \rho_{\mathcal{C}}(\bar{\tau}f)$ for all $f \in \text{FVL}[F]$. Let $X \in \mathcal{C}$ and let $T : F \rightarrow X$ be a contraction. There exists a contraction $S : E \rightarrow X$ so that $S\iota = T$. Suppose that $f = G(\delta_{y_1}, \dots, \delta_{y_n})$, where G is a lattice-linear expression and $y_1, \dots, y_n \in F$. We have

$$\widehat{S}(\bar{\tau}f) = \widehat{S}\widehat{\phi_{E\iota}}(f) = G(S\iota y_1, \dots, S\iota y_n) = G(Ty_1, \dots, Ty_n) = \widehat{T}f.$$

Thus, $\|\widehat{T}f\| \leq \rho_{\mathcal{C}}(\bar{\tau}f)$. Taking supremum over all such T shows that $\rho_{\mathcal{C}}(f) \leq \rho_{\mathcal{C}}(\bar{\tau}f)$. \square

Using the above results we may now characterize when $\text{FBL}^{\mathcal{C}}[F]$ embeds via $\bar{\tau}$ onto a complemented sublattice of $\text{FBL}^{\mathcal{C}}[E]$. This characterization should be contrasted with [28, Proposition 3.12] and our solution to the subspace problem in Theorem 5.8. Note, in particular, that if condition (1) in Theorem 5.4 holds for a class \mathcal{C} then it also holds for any class \mathcal{C}' such that $\mathcal{C}' \subseteq \mathcal{C}$.

Theorem 5.4. *Let F be a closed subspace of a Banach space E and let $\iota : F \rightarrow E$ be the inclusion map. Set $\bar{\tau} = \widehat{\phi_{E\iota}}$. The following are equivalent.*

- (1) *For any $X \in \bar{\mathcal{C}}$, any bounded linear operator $T : F \rightarrow X$ has a bounded linear extension $\tilde{T} : E \rightarrow X$ with $\|\tilde{T}\| = \|T\|$.*
- (2) *$\bar{\tau}$ is a lattice isometric embedding and there is a contractive lattice homomorphic projection P from $\text{FBL}^{\mathcal{C}}[E]$ onto $\bar{\tau}(\text{FBL}^{\mathcal{C}}[F])$.*

Proof. (1) \implies (2): By the above, $\bar{\tau}$ is a contraction. Furthermore, $\bar{\tau} = \widehat{\phi_E \bar{\iota}}$ is a lattice homomorphism. Since $\text{FBL}^{\mathcal{C}}[F] \in \bar{\mathcal{C}}$, the isometric embedding $\phi_F : F \rightarrow \text{FBL}^{\mathcal{C}}[F]$ has a bounded linear extension $\widetilde{\phi}_F : E \rightarrow \text{FBL}^{\mathcal{C}}[F]$ such that $\|\widetilde{\phi}_F\| = 1$. Therefore, $\widehat{\phi}_F : \text{FBL}^{\mathcal{C}}[E] \rightarrow \text{FBL}^{\mathcal{C}}[F]$ is a lattice homomorphism such that $\|\widehat{\phi}_F\| = \|\widetilde{\phi}_F\| = 1$ and $\widehat{\phi}_F \phi_E = \widetilde{\phi}_F$. For $y \in F$, it is easy to see that

$$\widehat{\phi}_F \bar{\iota} \delta_y = \widehat{\phi}_F \delta_{\iota y} = \widetilde{\phi}_F(\iota y) = \phi_F y = \delta_y.$$

Since $\widehat{\phi}_F \bar{\iota}$ is a lattice homomorphism, it follows that $\widehat{\phi}_F \bar{\iota}$ is the identity map on $\text{FBL}^{\mathcal{C}}[F]$. In particular,

$$\rho_{\mathcal{C}}(f) = \rho_{\mathcal{C}}(\widehat{\phi}_F \bar{\iota} f) \leq \rho_{\mathcal{C}}(\bar{\iota} f) \leq \rho_{\mathcal{C}}(f) \text{ for any } f \in \text{FBL}^{\mathcal{C}}[F].$$

Hence, $\bar{\tau}$ is a lattice isometric embedding. Furthermore, $P := \bar{\tau} \widehat{\phi}_F$ is a lattice homomorphic projection on $\text{FBL}^{\mathcal{C}}[E]$ such that $P \text{FBL}^{\mathcal{C}}[E] = \bar{\tau} \text{FBL}^{\mathcal{C}}[F]$ and $\|P\| = 1$.

(2) \implies (1): Let $X \in \bar{\mathcal{C}}$ and let $T : F \rightarrow X$ be a bounded linear operator. Then $\widehat{T} : \text{FBL}^{\mathcal{C}}[F] \rightarrow X$ is a lattice homomorphism such that $\|\widehat{T}\| = \|T\|$ and $\widehat{T} \phi_F = T$. Let $\widetilde{T} = \widehat{T}^{-1} P \phi_E : E \rightarrow X$. For any $y \in F$,

$$\widetilde{T} y = \widehat{T}^{-1} P \delta_{\iota y} = \widehat{T}^{-1} \delta_{\iota y} = \widehat{T} \delta_y = T y.$$

Hence, \widetilde{T} is a bounded linear extension of T . Clearly, $\|\widetilde{T}\| \leq \|\widehat{T}\| = \|T\|$. Hence, $\|\widetilde{T}\| = \|T\|$. \square

We conclude this preliminary subsection with another simple observation.

Proposition 5.5. *Assume that $\bar{\tau} : \text{FBL}^{\mathcal{C}}[F] \rightarrow \text{FBL}^{\mathcal{C}}[E]$ is a lattice isometric embedding. Let $X \in \mathcal{C}$ and let $T : F \rightarrow X$ be a bounded linear operator. The following are equivalent.*

- (1) *There exists an operator $S : E \rightarrow X$ such that $S \bar{\iota} = T$ and $\|S\| = \|T\|$.*
- (2) *There exists a lattice homomorphism $R : \text{FBL}^{\mathcal{C}}[E] \rightarrow X$ such that $R \bar{\tau} = \widehat{T}$ and $\|R\| = \|T\|$.*

Proof. (1) \implies (2): Suppose that S is as given. Then $\widehat{S} : \text{FBL}^{\mathcal{C}}[E] \rightarrow X$ is a lattice homomorphism such that $\|\widehat{S}\| = \|S\| = \|T\|$. If $f = G(\delta_{y_1}, \dots, \delta_{y_n}) \in \text{FVL}[F]$, then from the proof of Proposition 5.2, $\bar{\tau} f = G(\delta_{\iota y_1}, \dots, \delta_{\iota y_n})$. Thus,

$$\widehat{S} \bar{\tau} f = \widehat{S} G(\delta_{\iota y_1}, \dots, \delta_{\iota y_n}) = G(S \iota y_1, \dots, S \iota y_n) = G(T y_1, \dots, T y_n) = \widehat{T} f,$$

so (2) holds with $R = \widehat{S}$.

(2) \implies (1): Assume that R is as given. Let $S = R \phi_E : E \rightarrow X$. Then $\|S\| = \|R\| = \|T\|$. Moreover, if $y \in F$ then

$$S \iota y = R \phi_E \iota y = R \delta_{\iota y} = R \bar{\tau} \delta_y = \widehat{T} \delta_y = T y.$$

Thus, (1) holds. \square

5.1. Push-outs in \mathcal{C} . Recall that for objects A_0, A_1, A_2 and morphisms $\alpha_i : A_0 \rightarrow A_i$, $i = 1, 2$, a push-out diagram is an object $PO = PO(\alpha_1, \alpha_2)$ together with morphisms $\beta_i : A_i \rightarrow PO$, $i = 1, 2$, making the following diagram commutative

$$\begin{array}{ccc} A_1 & \xrightarrow{\beta_1} & PO \\ \alpha_1 \uparrow & & \uparrow \beta_2 \\ A_0 & \xrightarrow{\alpha_2} & A_2 \end{array}$$

and with the universal property that if $\beta'_i : A_i \rightarrow B$ are such that $\beta'_1\alpha_1 = \beta'_2\alpha_2$, then there is a unique $\gamma : PO \rightarrow B$ such that $\gamma\beta_i = \beta'_i$ for $i = 1, 2$, as follows:

$$\begin{array}{ccccc}
 & & & & B \\
 & & & \nearrow^{\beta'_1} & \\
 & & & \gamma & \\
 & & & \searrow_{\beta'_2} & \\
 A_1 & \xrightarrow{\beta_1} & PO & & \\
 \alpha_1 \uparrow & & \beta_2 \uparrow & & \\
 A_0 & \xrightarrow{\alpha_2} & A_2 & &
 \end{array}$$

In the category of Banach lattices and lattice homomorphisms, isometric push-outs were shown to exist in [8] – these are push-outs satisfying the extra condition that $\max\{\|\beta_1\|, \|\beta_2\|\} \leq 1$ and which guarantee the inequality $\|\gamma\| \leq \max\{\|\beta'_1\|, \|\beta'_2\|\}$ in the universal property. The construction of such push-outs can be adapted to Banach lattices within certain classes $\overline{\mathcal{C}}$, as follows.

Theorem 5.6. *Let \mathcal{C} be a class of Banach lattices such that $\overline{\mathcal{C}}$ is closed under lattice quotients. Given Banach lattices X_0, X_1, X_2 in $\overline{\mathcal{C}}$ and lattice homomorphisms $T_i : X_0 \rightarrow X_i$ for $i = 1, 2$, there is a Banach lattice $PO^{\overline{\mathcal{C}}}$ in $\overline{\mathcal{C}}$ and lattice homomorphisms S_1, S_2 so that the following is an isometric push-out diagram in $\overline{\mathcal{C}}$:*

$$\begin{array}{ccc}
 X_1 & \xrightarrow{S_1} & PO^{\overline{\mathcal{C}}} \\
 T_1 \uparrow & & S_2 \uparrow \\
 X_0 & \xrightarrow{T_2} & X_2
 \end{array}$$

Proof. Let $X_1 \oplus_1 X_2$ denote the direct sum equipped with the norm $\|(x_1, x_2)\| = \|x_1\| + \|x_2\|$ and let $j_i : X_i \rightarrow X_1 \oplus_1 X_2$ denote the canonical embedding for $i = 1, 2$. Let $\phi : X_1 \oplus_1 X_2 \rightarrow \text{FBL}^{\mathcal{C}}[X_1 \oplus_1 X_2]$ be the canonical embedding and Z be the (closed) ideal in $\text{FBL}^{\mathcal{C}}[X_1 \oplus_1 X_2]$ generated by the families $(\phi(j_1|x|) - |\phi(j_1x)|)_{x \in X_1}$, $(\phi(j_2|y|) - |\phi(j_2y)|)_{y \in X_2}$ and $(\phi j_1 T_1 z - \phi j_2 T_2 z)_{z \in X_0}$. Let

$$PO^{\overline{\mathcal{C}}} = \text{FBL}^{\mathcal{C}}[X_1 \oplus_1 X_2]/Z,$$

and let $S_i = q\phi j_i : X_i \rightarrow PO^{\overline{\mathcal{C}}}$, for $i = 1, 2$, where $q : \text{FBL}^{\mathcal{C}}[X_1 \oplus_1 X_2] \rightarrow PO^{\overline{\mathcal{C}}}$ denotes the canonical quotient map.

Since $\overline{\mathcal{C}}$ is closed under lattice quotients and $\text{FBL}^{\mathcal{C}}[X_1 \oplus_1 X_2]$ is in $\overline{\mathcal{C}}$, so is $PO^{\overline{\mathcal{C}}}$. The rest of the proof follows exactly as in [8, Theorem 4.3]. \square

Theorem 5.6 applies, in particular, when \mathcal{C} is the class of p -convex Banach lattices or the class of Banach lattices with an upper p -estimate (with constant one). Moreover, as in [8, Theorem 4.4], for these classes we have the following.

Theorem 5.7. *Let \mathcal{C} be the class of p -convex Banach lattices or Banach lattices with an upper p -estimate. Set $K_{\mathcal{C}} = 2^{1-\frac{1}{p}}$ if $\mathcal{C} = \mathcal{C}_p$ and $K_{\mathcal{C}} = 2^{1-\frac{1}{p}}\gamma_p$ if $\mathcal{C} = \mathcal{C}_{p,\infty}$, and let*

$$\begin{array}{ccc}
 X_1 & \xrightarrow{\tilde{T}_1} & PO^{\overline{\mathcal{C}}} \\
 T_1 \uparrow & & \tilde{T}_2 \uparrow \\
 X_0 & \xrightarrow{T_2} & X_2
 \end{array}$$

be an isometric push-out diagram in $\overline{\mathcal{C}}$. If $\|T_1\| \leq 1$ and the lower arrow T_2 is an isometric embedding then the upper arrow \tilde{T}_1 is a $K_{\mathcal{C}}$ -embedding.

Proof. Fix $x_1 \in X_1$. We aim to show that $\|x_1\| \leq K_{\mathcal{C}}\|\tilde{T}_1(x_1)\|$. The reverse inequality is trivially true since $\|\tilde{T}_1\| \leq 1$. Since we are dealing with lattice norms and lattice homomorphisms, we can suppose that x_1 is positive. By the push-out universal property, as given in Theorem 5.6, it is enough to find a commutative diagram in $\overline{\mathcal{C}}$

$$\begin{array}{ccc} X_1 & \xrightarrow{\hat{T}_1} & Z \\ T_1 \uparrow & & \hat{T}_2 \uparrow \\ X_0 & \xrightarrow{T_2} & X_2 \end{array}$$

such that $\|x_1\| \leq \|\hat{T}_1(x_1)\|$ and $\max\{\|\hat{T}_1\|, \|\hat{T}_2\|\} \leq K_{\mathcal{C}}$.

To this end, by [8, Theorem 4.4] we have a Banach lattice isometric push-out diagram

$$\begin{array}{ccc} X_1 & \xrightarrow{\tilde{T}_1} & PO \\ T_1 \uparrow & & \tilde{T}_2 \uparrow \\ X_0 & \xrightarrow{T_2} & X_2 \end{array}$$

such that \tilde{T}_1 is an isometric embedding. Hence, by [8, Lemma 3.3], we can pick a positive σ -finite element $x^* \in PO^*$ of norm one such that $x^*(|\tilde{T}_1 x_1|) = \|x_1\|$.

The functional x^* induces a lattice semi-norm on PO given by $\|z\|_{x^*} = x^*(|z|)$. After making a quotient by the elements of norm 0, this becomes a norm that satisfies $\| |x| + |y| \|_{x^*} = \|x\|_{x^*} + \|y\|_{x^*}$ for all x, y . This identity extends to the completion of this normed lattice, which, by Kakutani's representation theorem [22, Theorem 1.b.2], is lattice isometric to a Banach lattice of the form $L^1(\nu_{x^*})$. The formal identity induces a homomorphism $\psi_{x^*} : PO \rightarrow L^1(\nu_{x^*})$ of norm one.

For $i = 1, 2$, let $A_i = \psi_{x^*}(\tilde{T}_i(B_{X_i}))$ and set $A = A_1 \cup A_2 \subseteq L^1(\nu_{x^*})$. We will distinguish the following cases:

- (1) Let $\mathcal{C} = \mathcal{C}_p$ denote the class of p -convex Banach lattices (with constant 1). Since the X_i are p -convex and the $\Psi_{x^*}\tilde{T}_i$ are contractive lattice homomorphisms we have that for all finitely supported sequences $(\alpha_i)_{i \in I}$ of real numbers and $(f_i)_{i \in I} \subseteq A$,

$$\left\| \left(\sum |\alpha_i f_i|^p \right)^{\frac{1}{p}} \right\|_1 \leq \left\| \left(\sum_{f_i \in A_1} |\alpha_i f_i|^p \right)^{\frac{1}{p}} \right\|_1 + \left\| \left(\sum_{f_i \in A_2} |\alpha_i f_i|^p \right)^{\frac{1}{p}} \right\|_1 \leq 2^{1-\frac{1}{p}} \left(\sum |\alpha_i|^p \right)^{\frac{1}{p}}.$$

Hence, by Theorem 3.1, there is $g \in L^1(\nu_{x^*})_+$ with $\|g\|_1 \leq 1$ and lattice homomorphisms $\hat{T}_i : X_i \rightarrow L^p(g\nu_{x^*})$ that make the following diagram commutative

$$\begin{array}{ccccc} X_i & \xrightarrow{\tilde{T}_i} & PO & \xrightarrow{\psi_{x^*}} & L^1(\nu_{x^*}) \\ & \searrow \hat{T}_i & & \nearrow M_g & \\ & & L^p(g\nu_{x^*}) & & \end{array}$$

with $\|\widehat{T}_i\| \leq 2^{1-\frac{1}{p}} = K_{\mathcal{C}_p}$ and M_g the operator of multiplication by g . It follows that

$$\begin{array}{ccc} X_1 & \xrightarrow{\widehat{T}_1} & L^p(g\nu_{x^*}) \\ T_1 \uparrow & & \widehat{T}_2 \uparrow \\ X_0 & \xrightarrow{T_2} & X_2 \end{array}$$

is a commutative diagram in \mathcal{C}_p and

$$\|x_1\| = x^*|\widetilde{T}_1 x_1| = \|\psi_{x^*}\widetilde{T}_1 x_1\|_{L^1(\nu_{x^*})} \leq \|\widehat{T}_1 x_1\|_{L^p(g\nu_{x^*})}.$$

- (2) Let $\mathcal{C} = \mathcal{C}_{p,\infty}$ denote the class of Banach lattices satisfying an upper p -estimate (with constant 1). Since the X_i satisfy an upper p -estimate and the $\Psi_{x^*}\widehat{T}_i$ are contractive lattice homomorphisms we have that for all finitely supported sequences $(\alpha_i)_{i \in I}$ of real numbers and $(f_i)_{i \in I} \subseteq A$,

$$\left\| \bigvee |\alpha_i f_i| \right\|_1 \leq \left\| \bigvee_{f_i \in A_1} |\alpha_i f_i| \right\|_1 + \left\| \bigvee_{f_i \in A_2} |\alpha_i f_i| \right\|_1 \leq 2^{1-\frac{1}{p}} \left(\sum |\alpha_i|^p \right)^{\frac{1}{p}}.$$

Hence, by Theorem 3.2, there is $g \in L^1(\nu_{x^*})_+$ with $\|g\|_1 \leq 1$ and lattice homomorphisms $\widehat{T}_i : X_i \rightarrow L^{p,\infty}(g\nu_{x^*})$ that make the following diagram commutative

$$\begin{array}{ccccc} X_i & \xrightarrow{\widetilde{T}_i} & PO & \xrightarrow{\psi_{x^*}} & L^1(\nu_{x^*}) \\ & \searrow \widehat{T}_i & & \nearrow M_g & \\ & & L^{p,\infty}(g\nu_{x^*}) & & \end{array}$$

with $\|\widehat{T}_i\| \leq 2^{1-\frac{1}{p}}\gamma_p = K_{\mathcal{C}_{p,\infty}}$ and M_g the operator of multiplication by g . It follows that

$$\begin{array}{ccc} X_1 & \xrightarrow{\widehat{T}_1} & L^{p,\infty}(g\nu_{x^*}) \\ T_1 \uparrow & & \widehat{T}_2 \uparrow \\ X_0 & \xrightarrow{T_2} & X_2 \end{array}$$

is a commutative diagram in $\mathcal{C}_{p,\infty}$ and

$$\|x_1\| = x^*|\widetilde{T}_1 x_1| = \|\psi_{x^*}\widetilde{T}_1 x_1\|_{L^1(\nu_{x^*})} \leq \|\widehat{T}_1 x_1\|_{L^{p,\infty}(g\nu_{x^*})}.$$

□

As a consequence of the above, we get our desired solution to the subspace problem.

Theorem 5.8. *Suppose that $\iota : F \rightarrow E$ is an isometric embedding and let \mathcal{C} denote either the class of p -convex Banach lattices or the class of Banach lattices with upper p -estimates. The following are equivalent.*

- (1) $\bar{\iota} : \text{FBL}^{\mathcal{C}}[F] \rightarrow \text{FBL}^{\mathcal{C}}[E]$ is a c_1 -lattice embedding.
- (2) For every operator $T : F \rightarrow X$ with X in \mathcal{C} , there is Y in \mathcal{C} , a (norm one) $K_{\mathcal{C}}$ -lattice embedding $j : X \rightarrow Y$ and $S : E \rightarrow Y$ so that $jT = S\iota$ and $\|S\| \leq c_2\|T\|$.

Here, $c_2 \leq c_1 \leq K_{\mathcal{C}}c_2$.

Proof. (1) \implies (2): Let $X \in \mathcal{C}$ and $T : F \rightarrow X$ an operator. By Theorem 5.6, let us consider the isometric push-out diagram in $\overline{\mathcal{C}}$ for $X_0 = \overline{\tau}(\text{FBL}^{\mathcal{C}}(F))$, $X_1 = X$, $X_2 = \text{FBL}^{\mathcal{C}}(E)$, $T_1 = \frac{\widehat{T}\overline{\tau}^{-1}}{\|\widehat{T}\overline{\tau}^{-1}\|} : X_0 \rightarrow X_1$ and T_2 the formal identity from X_0 to X_2 , where $\widehat{T} : \text{FBL}^{\mathcal{C}}[F] \rightarrow X$ denotes the lattice homomorphism extending T .

$$\begin{array}{ccccc}
& & X & \xrightarrow{S_1} & PO\overline{\mathcal{C}} \\
& \nearrow \frac{\widehat{T}}{\|\widehat{T}\overline{\tau}^{-1}\|} & \uparrow T_1 & & \uparrow S_2 \\
\text{FBL}^{\mathcal{C}}[F] & \xrightarrow{\overline{\tau}} & \overline{\tau}(\text{FBL}^{\mathcal{C}}(F)) & \xrightarrow{T_2} & \text{FBL}^{\mathcal{C}}[E] \\
\uparrow \phi_F & & & & \uparrow \phi_E \\
F & \xrightarrow{\iota} & & & E
\end{array}$$

Let us write $Y = PO\overline{\mathcal{C}}$ and $j = S_1$. Since T_2 is an isometric embedding, we can apply Theorem 5.7 to obtain that j is a norm one $K_{\mathcal{C}}$ -embedding. On the other hand, let $S = \|\widehat{T}\overline{\tau}^{-1}\|S_2\phi_E : E \rightarrow Y$. It follows that

$$jT = j\widehat{T}\phi_F = \|\widehat{T}\overline{\tau}^{-1}\|jT_1\overline{\tau}\phi_F = \|\widehat{T}\overline{\tau}^{-1}\|S_2T_2\overline{\tau}\phi_F = \|\widehat{T}\overline{\tau}^{-1}\|S_2\phi_E\iota = S\iota,$$

and $\|S\| \leq \|\widehat{T}\|\|\overline{\tau}^{-1}\| \leq c_1\|T\|$, so $c_2 \leq c_1$.

(2) \implies (1): First, note that $\|\overline{\tau}\| = 1$. Now, take $f \in \text{FVL}[F]$ and a contraction $T : F \rightarrow X$, X in \mathcal{C} . By the hypothesis, there exists Y in \mathcal{C} , a norm one $K_{\mathcal{C}}$ -lattice embedding $j : X \rightarrow Y$ and $S : E \rightarrow Y$ so that $jT = S\iota$ and $\|S\| \leq c_2$. It follows that $j\widehat{T} = \widehat{S}\overline{\tau}$, so

$$\|\widehat{T}f\|_X = \|j^{-1}\widehat{S}\overline{\tau}f\|_X \leq \|j^{-1}\|\|S\|\rho_{\mathcal{C}}(\overline{\tau}f) \leq K_{\mathcal{C}}c_2\rho_{\mathcal{C}}(\overline{\tau}f)$$

and $\overline{\tau}$ is a $K_{\mathcal{C}}c_2$ -embedding. \square

5.2. Recovering the POE- p via injectivity. Recall that a Banach lattice X is *injective* if for every Banach lattice Z , every closed linear sublattice Y of Z and every positive linear operator $T : Y \rightarrow X$ there is a positive linear extension $\widehat{T} : Z \rightarrow X$ with $\|\widehat{T}\| = \|T\|$. Equivalently, X is injective if whenever X lattice isometrically embeds into a Banach lattice Y there is a positive contractive projection from Y onto X . Note that this notion of injectivity corresponds to the category of Banach lattices with positive linear maps, which is considerably different from the category of Banach lattices with lattice homomorphisms (where in fact there are no injective objects [8]). The study of injective Banach lattices is classical [12, 20, 23]; however, it seems that not much is known on injectivity (or projectivity) in subcategories of Banach lattices. Here, we study the injectivity of ℓ^p in the class of p -convex Banach lattices, with an aim towards showing that Theorem 5.8 together with Proposition 5.3 recover [28, Theorem 3.7], up to constants. Although the injectivity of L^p spaces in the p -convex category is implicit in [31], we provide an independent argument in order to give an alternative proof of [28, Theorem 3.7]. See also [17, Corollary 4.13] for a related result.

Theorem 5.9. *Let X be a p -convex Banach lattice and suppose that (x_n) is a disjoint positive sequence in X equivalent to the ℓ^p -basis. Then $[(x_n)]$ is the image of a positive projection on X .*

Proof. We may assume that X is a closed sublattice of $Z = (\bigoplus_{\mu \in \Gamma} L^p(\mu))_\infty$ by Proposition 3.3. It suffices to show that $[(x_n)]$ is the range of a positive projection on Z . Assume that $c\|(a_n)\|_p < \|\sum a_n x_n\| \leq \|(a_n)\|_p$ for any non-zero $(a_n) \in \ell^p$. Let $m \in \mathbb{N}$. There exists a finite set Γ_m , $k \in \mathbb{N}$, and a positive contraction $T : (\bigoplus_{\mu \in \Gamma} L^p(\mu))_\infty \rightarrow (\ell^p(k))_\infty^m$ so that $(Tx_n)_{n=1}^m$ is a positive disjoint sequence satisfying $c\|(a_n)_{n=1}^m\|_p < \|\sum_{n=1}^m a_n Tx_n\| \leq \|(a_n)_{n=1}^m\|_p$ for any non-zero $(a_n)_{n=1}^m \in \ell^p(m)$. Write $Tx_n = (y_n(\gamma))_{\gamma \in \Gamma_m}$ with $y_n(\gamma) \in \ell^p(k)$. Define $S : \ell^1(m) \rightarrow \ell^\infty(\Gamma_m)$ by

$$S(b_1, \dots, b_m) = \left(\sum_{n=1}^m b_n \|y_n(\gamma)\|^p \right)_{\gamma \in \Gamma_m}.$$

Since $(y_n(\gamma))_{n=1}^m$ is a disjoint sequence for each γ , if $b = (b_1, \dots, b_m) \geq 0$, then

$$\|Sb\| = \max_{\gamma \in \Gamma_m} \left\| \sum_{n=1}^m b_n^{\frac{1}{p}} y_n(\gamma) \right\|_p^p = \left\| \sum_{n=1}^m b_n^{\frac{1}{p}} Tx_n \right\|^p \geq c^p \|(b_n^{\frac{1}{p}})_{n=1}^m\|_p^p = c^p \|b\|_1.$$

We claim that $B_{\ell^\infty(m)}^+ \subseteq c^{-p} \text{co so}\{S^* e_\gamma^* : \gamma \in \Gamma_m\}$. Here, for a subset A of a Banach lattice X , so $A = \bigcup_{a \in A} [-|a|, |a|]$ denotes the solid hull of A and, as usual, $\text{co}(A)$ denotes the convex hull. To prove the claim, note that if $0 \leq y^* \notin c^{-p} \text{co so}\{S^* e_\gamma^* : \gamma \in \Gamma_m\}$ then by [29, Proposition 3.1] there exists $y \in \ell^1(m)_+$ so that

$$\|y^*\| \|y\| \geq y^*(y) > c^{-p} \max_{\gamma \in \Gamma_m} (S^* e_\gamma^*)(y) = c^{-p} \|Sy\| \geq \|y\|.$$

Hence, $\|y^*\| > 1$. This completes the proof of the claim.

By the claim we can find a convex combination $\sum_{\gamma \in \Gamma_m} \alpha_\gamma S^* e_\gamma^* \geq c^p \overbrace{(1, \dots, 1)}^m$. Choose $y_n^*(\gamma) \in \ell^{p'}(k)_+$ so that $\text{supp } y_n^*(\gamma) = \text{supp } y_n(\gamma) =: I_n$, $\|y_n^*(\gamma)\|_{p'} = 1$ and $\langle y_n(\gamma), y_n^*(\gamma) \rangle = \|y_n(\gamma)\|_p$. If $x \in (\bigoplus_{\mu \in \Gamma} L^p(\mu))_\infty$ and $Tx = (y(\gamma))_{\gamma \in \Gamma_m}$, let

$$z_n^*(x) = \sum_{\gamma \in \Gamma_m} \alpha_\gamma \|y_n(\gamma)\|_p^{p-1} \langle y(\gamma), y_n^*(\gamma) \rangle.$$

It is easy to see that z_n^* is a positive linear functional and for any $x \in (\bigoplus_{\mu \in \Gamma} L^p(\mu))_\infty$,

$$\begin{aligned} \|(z_n^*(x))_{n=1}^m\|_p &\leq \sum_{\gamma \in \Gamma_m} \alpha_\gamma \left\| (\|y_n(\gamma)\|_p^{p-1} \|y(\gamma)\|_p \chi_{I_n})_{n=1}^m \right\|_p \\ &\leq \sup_{\gamma \in \Gamma_m} \sup_n \|y_n(\gamma)\|_p^{p-1} \left\| \sum_{n=1}^m y(\gamma) \chi_{I_n} \right\|_p \\ &\leq \sup_{\gamma \in \Gamma_m} \sup_n \|y_n(\gamma)\|_p^{p-1} \|y(\gamma)\|_p \\ &\leq \sup_n \|Tx_n\|_p^{p-1} \cdot \|Tx\| \leq \|Tx\| \leq \|x\|. \end{aligned}$$

On the other hand, $z_l^*(x_n) = 0$ if $l \neq n$ and

$$z_n^*(x_n) = \sum_{\gamma \in \Gamma_m} \alpha_\gamma \|y_n(\gamma)\|_p^{p-1} \langle y_n(\gamma), y_n^*(\gamma) \rangle = \sum_{\gamma \in \Gamma_m} \alpha_\gamma \|y_n(\gamma)\|_p^p \geq c^p,$$

since $\sum_{\gamma \in \Gamma_m} \alpha_\gamma \|y_n(\gamma)\|_p^p$ is the n -th coordinate of the sum $\sum_{\gamma \in \Gamma_m} \alpha_\gamma S^* e_\gamma^*$. Therefore, the map P_m on $(\bigoplus_{\mu \in \Gamma} L^p(\mu))_\infty$ given by

$$P_m x = \sum_{n=1}^m \frac{z_n^*(x)}{z_n^*(x_n)} x_n$$

is a positive projection from $(\bigoplus_{\mu \in \Gamma} L^p(\mu))_\infty$ onto $[(x_n)_{n=1}^m]$ such that $\|P_m\| \leq c^{-p}$. Finally, let \mathcal{U} be a free ultrafilter on \mathbb{N} . Regard $[(x_n)_{n=1}^\infty]$ as a dual Banach lattice (either the dual of an isomorph of $\ell^{p'}$ if $1 < p \leq \infty$ or the dual of an isomorph of c_0 if $p = 1$). The operator $P : (\bigoplus_{\mu \in \Gamma} L^p(\mu))_\infty \rightarrow [(x_n)]$ given by $Px = w^*\text{-lim}_{m \rightarrow \mathcal{U}} P_m x$ is a positive projection onto $[(x_n)]$. \square

As a corollary of Theorems 5.8 and 5.9, we recover the solution to the subspace problem for $\text{FBL}^{(p)}$ given in [28, Theorem 3.7].

Corollary 5.10. *Let $\iota : F \hookrightarrow E$ be an isometric embedding and $1 \leq p < \infty$. The following are equivalent.*

- (1) $\tau : \text{FBL}^{(p)}[F] \rightarrow \text{FBL}^{(p)}[E]$ is a c_1 -lattice embedding.
- (2) For every operator $T : F \rightarrow \ell^p$ there is $R : E \rightarrow \ell^p$ so that $T = R\iota$ and $\|R\| \leq c_2 \|T\|$.

Here, $c_1 \leq c_2 \leq 2^{p-\frac{1}{p}} c_1$.

Proof. We use the identification $\text{FBL}^{(p)} = \text{FBL}^{\mathcal{C}}$ where \mathcal{C} denotes the class of p -convex Banach lattices and invoke Theorem 5.8. The implication (2) \Rightarrow (1) is clear. For the converse, we get that there exist a p -convex Banach lattice Y , a $2^{1-\frac{1}{p}}$ -lattice embedding $j : \ell^p \rightarrow Y$ and $S : E \rightarrow Y$ such that $jT = S\iota$ with $\|S\| \leq c_1 \|T\|$. By Theorem 5.9, there is a (positive) projection $P : Y \rightarrow j(\ell^p) \subseteq Y$ with $\|P\| \leq (2^{1-\frac{1}{p}})^p$ (as j is a $2^{1-\frac{1}{p}}$ -embedding). Thus, we can take $R = j^{-1}PS$. \square

5.3. $\ell^{p,\infty}$ is not injective for Banach lattices with upper p -estimates. We now show that for any $1 < p < \infty$, $\ell^{p,\infty}$ is not injective in the class of Banach lattices with upper p -estimates.

Theorem 5.11. *For any $1 < p < \infty$ there is a Banach lattice X with an upper p -estimate and an uncomplemented closed sublattice Y of X that is lattice isomorphic to $\ell^{p,\infty}$.*

Proof. We begin with some notation. Let X be the Banach lattice $(\bigoplus_{\mathbb{N}} \ell^{p,\infty})_\infty$. Clearly, X satisfies an upper p -estimate with constant 1. We may express any $x \in X$ as $x = (x(j))_{j=1}^\infty = (x(i, j))_{i,j=1}^\infty$, where $(x(j))$ is a bounded sequence in $\ell^{p,\infty}$ and $x(j) = (x(i, j))_{i=1}^\infty$ for each j . Given $m \in \mathbb{N}$, define P_m on X by

$$(P_m x)(i, j) = \begin{cases} x(m, j) & \text{if } i = m, \\ 0 & \text{if } i \neq m. \end{cases}$$

Clearly, P_m is a band projection on X and the projection band $P_m X$ is lattice isometric to ℓ^∞ via the map

$$L_m : \ell^\infty \rightarrow P_m X, (L_m y)(i, j) = \begin{cases} y(j) & \text{if } i = m, \text{ where } y = (y(j))_{j=1}^\infty, \\ 0 & \text{otherwise.} \end{cases}$$

Let \mathcal{S} be the Schreier family, i.e., all subsets I of \mathbb{N} so that $|I| \leq \min I$. Since \mathcal{S} is countable, we can list its elements in a sequence $(S_j)_{j=1}^\infty$. Given $a = (a_i) \in \ell^{p,\infty}$, let

$$x_a(i, j) = \begin{cases} a_i & \text{if } i \in S_j, \\ 0 & \text{otherwise.} \end{cases}$$

Denote the sequence of coordinate unit vectors in $\ell^{p,\infty}$ by (e_i) . For any j ,

$$\|(x_a(i, j))_{i=1}^\infty\|_{p,\infty} = \left\| \sum_{i \in S_j} a_i e_i \right\|_{p,\infty} \leq \|(a_i)\|_{p,\infty}.$$

Hence, $T : \ell^{p,\infty} \rightarrow X$, $a \mapsto x_a$ is a bounded linear map which is clearly a lattice homomorphism. On the other hand, if $\|a\|_{p,\infty} > 1$ then there exists a non-empty finite set L in \mathbb{N} so that $|a_i| \geq (p'|L|^{\frac{1}{p}})^{-1}$ for all $i \in L$. Let $S_j \in \mathcal{S}$ be such that $S_j \subseteq L$ and $|S_j| \geq |L|/2$. Then,

$$\|x_a\| \geq \left\| \sum_{i \in S_j} a_i e_i \right\|_{p,\infty} \geq (p'|L|^{\frac{1}{p}})^{-1} \left\| \sum_{i \in S_j} e_i \right\|_{p,\infty} = \frac{1}{p'} \left(\frac{|S_j|}{|L|} \right)^{\frac{1}{p'}} \geq \frac{1}{p'2^{\frac{1}{p'}}}.$$

This shows that T is a lattice isomorphism from $\ell^{p,\infty}$ into X .

Let $Y = T\ell^{p,\infty}$. We will show that there is no bounded projection from X onto Y . Assume on the contrary that there is a projection P from X onto Y . Then there is a sequence (x_i^*) in X^* so that $(x_i^*(x))_i \in \ell^{p,\infty}$ for any $x \in X$ and $x_i^*(Ta) = a_i$ for any i and $a = (a_k) \in \ell^{p,\infty}$. It follows readily that $(x_i^*)_i \subseteq X^*$ is equivalent to the unit vector basis of $\ell^{p',1}$. In particular, it is an unconditional basic sequence in a Banach lattice that is q -concave for some $q < \infty$. Since (P_i^*) is a disjoint sequence of band projections on X^* , the sequence $(P_i^*x_i^*)$ is disjoint and hence also unconditional in X^* . By [22, Theorem 1.d.6(i)], for any finitely nonzero real sequence (b_i) ,

$$\|(b_i)\|_{p',1} \sim \left\| \sum b_i x_i^* \right\| \sim \left\| \sqrt{\sum |b_i x_i^*|^2} \right\| \geq \left\| \sqrt{\sum |b_i P_i^* x_i^*|^2} \right\| \sim \left\| \sum b_i P_i^* x_i^* \right\|.$$

Thus, the linear map $Q : X \rightarrow \ell^{p,\infty}$ defined by $Qx = (P_i^*x_i^*(x))_i$ is bounded. On the other hand, observe that $Te_k \in P_k X$. Therefore,

$$(5.1) \quad \langle Te_k, P_i^* x_i^* \rangle = \begin{cases} \langle Te_i, x_i^* \rangle = 1 & \text{if } k = i, \\ 0 & \text{otherwise.} \end{cases}$$

Hence, $QT e_i = e_i$ for any $i \in \mathbb{N}$.

Regard L_i as a lattice isometric embedding from ℓ^∞ into X . Define $y_i^* = L_i^* P_i^* x_i^*$ in $(\ell^\infty)^*$. Let I be a finite subset of \mathbb{N} . Suppose that $(u_i)_{i \in I}$ is a sequence in ℓ^∞ so that $\|(\sum_{i \in I} |u_i|^p)^{\frac{1}{p}}\|_\infty \leq 1$. Set $u = \sum_{i \in I} L_i u_i \in X$. Then $\|u\| \leq \|(\sum_{i \in I} |u_i|^p)^{\frac{1}{p}}\|_\infty \leq 1$. Thus,

$$\begin{aligned} \sum_{i \in I} y_i^*(u_i) &= \sum_{i \in I} \langle u_i, L_i^* P_i^* x_i^* \rangle = \sum_{i \in I} (P_i^* x_i^*)(u) \\ &\leq \|(P_i^* x_i^*)(u)\|_{p,\infty} \cdot |I|^{\frac{1}{p'}} = \|Qu\| \cdot |I|^{\frac{1}{p'}} \lesssim |I|^{\frac{1}{p'}}. \end{aligned}$$

Making use of the duality described on [22, p. 47], we conclude that there is an absolute constant $C < \infty$ so that

$$(5.2) \quad \left\| \left(\sum_{i \in I} |y_i^*|^{p'} \right)^{\frac{1}{p'}} \right\| \leq C |I|^{\frac{1}{p'}}$$

for any finite subset I of \mathbb{N} . Since $(\ell^\infty)^*$ is an AL -space, the bounded sequence (y_i^*) has a subsequence $(y_{i_k}^*)$ that has a splitting

$$y_{i_k}^* = u_k^* + v_k^*, \quad |u_k^*| \wedge |v_k^*| = 0, \quad (u_k^*) \text{ is almost order bounded, } (v_k^*) \text{ is disjoint.}$$

See, e.g., [41]. Since $|v_k^*| \leq |y_{i_k}^*|$ for all k , it follows from (5.2) that $\|v_k^*\| \rightarrow 0$. Thus, the whole sequence $(y_{i_k}^*)$ is almost order bounded in $(\ell^\infty)^*$ and hence it is relatively weakly compact.

Since $x_i := Te_i \in P_i X$ and $L_i : \ell^\infty \rightarrow P_i X$ is a surjective lattice isometry, we may find a sequence (u_k) in ℓ^∞ so that $L_{i_k} u_k = x_{i_k}$ for each k . If we write $u_k = (u_k(j))_{j=1}^\infty$ then $u_k(j) = 1$ if $k \in S_j$ and 0 otherwise. Hence, for any finitely supported real sequence (a_k) ,

$$\left\| \sum a_k u_k \right\|_\infty = \sup_j \left| \sum_{k \in S_j} a_k \right|.$$

The expression on the right is the norm of the Schreier space. The preceding equation says that (u_k) is equivalent to the unit vector basis of the Schreier space. Thus, (u_k) is a weakly null sequence (in ℓ^∞). Since ℓ^∞ has the Dunford-Pettis property, the weakly null sequence (u_k) converges uniformly to 0 on the relatively weakly compact set $(y_{i_k}^*)$. In particular,

$$0 = \lim y_{i_k}^*(u_k) = \lim \langle u_k, L_{i_k}^* P_{i_k}^* x_{i_k}^* \rangle = \lim \langle x_{i_k}, P_{i_k}^* x_{i_k}^* \rangle.$$

However,

$$e_{i_k} = QTe_{i_k} = Qx_{i_k} = (P_j^* x_j^*(x_{i_k}))_j.$$

Taking the i_k -th component yields that $\langle x_{i_k}, P_{i_k}^* x_{i_k}^* \rangle = 1$ for any k , which is the desired contradiction. \square

A finite dimensional version of Theorem 5.11 can be achieved via a gluing argument. We begin with some preliminaries.

By the above, the map $T : \ell^{p,\infty} \rightarrow X := (\bigoplus_{\mathbb{N}} \ell^{p,\infty})_\infty$ given by $Ta = (a\chi_{S_j})_{j=1}^\infty$ where $a = (a_i) \in \ell^{p,\infty}$ and (S_j) is the sequence of Schreier sets is a lattice isomorphism so that $T(\ell^{p,\infty})$ is not complemented in X . For each $m \in \mathbb{N}$, regard $\ell^{p,\infty}(m)$ as the subspace $[(e_i)_{i=1}^m]$ in $\ell^{p,\infty}$. Define $T_m : \ell^{p,\infty}(m) \rightarrow X$ by $T_m = T|_{\ell^{p,\infty}(m)}$. Clearly, there is a constant C so that every T_m is a C -lattice embedding. If (E_m) , (F_m) are sequences of Banach spaces and $U_m : E_m \rightarrow F_m$ is a bounded linear operator for each m so that $\sup_m \|U_m\| < \infty$, we let $\bigoplus U_m : (\bigoplus E_m)_\infty \rightarrow (\bigoplus F_m)_\infty$ be the bounded linear operator defined by $(\bigoplus U_m)((u_m)_{m=1}^\infty) = (U_m u_m)_{m=1}^\infty$.

Corollary 5.12. *Let the notation be as above. Suppose that for each m , π_m is a projection from X onto $Y_m := T_m(\ell^{p,\infty}(m))$. Then $\sup_m \|\pi_m\| = \infty$.*

Proof. Assume, on the contrary, that there is such a sequence (π_m) so that $\sup \|\pi_m\| < \infty$. First of all, it is clear that the composition

$$\left(\bigoplus T_m^{-1} \right) \left(\bigoplus \pi_m \right) \left(\bigoplus T_m \right) = \bigoplus (T_m^{-1} \pi_m T_m)$$

is the identity operator on $(\bigoplus \ell^{p,\infty}(m))_\infty$. Let \mathcal{U} be a free ultrafilter on \mathbb{N} . If $b = (b_m) \in (\bigoplus \ell^{p,\infty}(m))_\infty$, $b_m = (b_m(j))_{j=1}^m$, let $Qb = (\lim_{m \rightarrow \mathcal{U}} b_m(j))_{j=1}^\infty$. It is clear that $Qb \in \ell^{p,\infty}$ and $Q : (\bigoplus \ell^{p,\infty}(m))_\infty \rightarrow \ell^{p,\infty}$ is a contraction. Define $J : \ell^{p,\infty} \rightarrow (\bigoplus \ell^{p,\infty}(m))_\infty$ by $Ja = (a\chi_{[1,m]})_{m=1}^\infty$. Then J is a lattice isometric embedding and $Q \bigoplus (T_m^{-1} \pi_m T_m) J = QJ$ is the identity map on $\ell^{p,\infty}$. Finally, define one more map $V : X = (\bigoplus \ell^{p,\infty})_\infty \rightarrow (\bigoplus X)_\infty$ as follows: for $c = (c_j)_{j=1}^\infty \in (\bigoplus \ell^{p,\infty})_\infty$, let

$$Vc = ((c_j \chi_{[1,m]})_{j=1}^\infty)_{m=1}^\infty.$$

Obviously, V is a contraction. Moreover, for any $a \in \ell^{p,\infty}$,

$$VTa = V((a\chi_{S_j})_{j=1}^\infty) = ((a\chi_{S_j} \chi_{[1,m]})_{j=1}^\infty)_{m=1}^\infty = (T_m(a\chi_{[1,m]}))_{m=1}^\infty = \left(\bigoplus T_m \right) Ja.$$

In summary, we have the commutative diagram

$$\begin{array}{ccccc} \ell^{p,\infty} & \xrightarrow{J} & (\bigoplus \ell^{p,\infty}(m))_\infty & \xrightarrow{\bigoplus T_m} & (\bigoplus X)_\infty \\ & \searrow T & & & \nearrow V \\ & & X & & \end{array}$$

and

$$\ell^{p,\infty} \xrightarrow{(\bigoplus T_m)^J} (\bigoplus X)_\infty \xrightarrow{Q(\bigoplus T_m^{-1}\pi_m)} \ell^{p,\infty}$$

is the identity map. Thus, the map $R := TQ(\bigoplus T_m^{-1}\pi_m)V : X \rightarrow X$ is a projection on X so that $RT = T$. Hence, $\text{range } R = T(\ell^{p,\infty})$, contrary to what we previously established for T . \square

ACKNOWLEDGEMENTS

The research of P. Tradacete and E. García-Sánchez is partially supported by the grants PID2020-116398GB-I00 and CEX2019-000904-S funded by MCIN/AEI/10.13039/501100011033. E. García-Sánchez is also partially supported by “European Union NextGenerationEU/PRT” and FPI grant CEX2019-000904-S-21-3 funded by MCIN/AEI/10.13039/501100011033. P. Tradacete is also supported by a 2022 Leonardo Grant for Researchers and Cultural Creators, BBVA Foundation.

We thank Timur Oikhberg for interesting discussions related to this work.

REFERENCES

- [1] Fernando Albiac and Nigel J. Kalton. *Topics in Banach space theory*, volume 233 of *Graduate Texts in Mathematics*. Springer, second edition, 2016.
- [2] Charalambos D. Aliprantis and Owen Burkinshaw. *Positive operators*. Springer, Dordrecht, 2006. Reprint of the 1985 original.
- [3] Antonio Avilés, Gonzalo Martínez-Cervantes, José Rodríguez, and Pedro Tradacete. A Godefroy-Kalton principle for free Banach lattices. *Israel J. Math.*, 247(1):433–458, 2022.
- [4] Antonio Avilés, Gonzalo Martínez-Cervantes, and José David Rodríguez Abellán. On projective Banach lattices of the form $C(K)$ and $FBL[E]$. *J. Math. Anal. Appl.*, 489(1):124129, 11, 2020.
- [5] Antonio Avilés, Gonzalo Martínez-Cervantes, and José David Rodríguez Abellán. On the Banach lattice c_0 . *Rev. Mat. Complut.*, 34(1):203–213, 2021.
- [6] Antonio Avilés, Grzegorz Plebanek, and José David Rodríguez Abellán. Chain conditions in free Banach lattices. *J. Math. Anal. Appl.*, 465(2):1223–1229, 2018.
- [7] Antonio Avilés, José Rodríguez, and Pedro Tradacete. The free Banach lattice generated by a Banach space. *J. Funct. Anal.*, 274(10):2955–2977, 2018.
- [8] Antonio Avilés and Pedro Tradacete. Amalgamation and injectivity in Banach lattices. *Int. Math. Res. Not. IMRN*, (2):956–997, 2023.
- [9] Colin Bennett and Robert Sharpley. *Interpolation of operators*, volume 129 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1988.
- [10] Eugene Bilokopytov. Order continuity and regularity on vector lattices and on lattices of continuous functions. *Positivity*, 27(4):Paper No. 52, 29, 2023.
- [11] Stan Byrd. Factoring operators satisfying p -estimates. *Trans. Amer. Math. Soc.*, 310(2):567–582, 1988.
- [12] Donald I. Cartwright. Extensions of positive operators between Banach lattices. *Mem. Amer. Math. Soc.*, 3(164):iv+48, 1975.
- [13] Si-Kit Chung and Denny Leung. On order continuous norms. *Proc. Amer. Math. Soc.*, 128(7):1971–1974, 2000.
- [14] Sheldon Dantas, Gonzalo Martínez-Cervantes, José David Rodríguez Abellán, and Abraham Rueda Zoca. Norm-attaining lattice homomorphisms. *Rev. Mat. Iberoam.*, 38(3):981–1002, 2022.

- [15] Ben de Pagter and Anthony W. Wickstead. Free and projective Banach lattices. *Proc. Roy. Soc. Edinburgh Sect. A*, 145(1):105–143, 2015.
- [16] Tadeusz Figiel. Uniformly convex norms on Banach lattices. *Studia Math.*, 68(3):215–247, 1980.
- [17] Julio Flores, Francisco L. Hernández, Eugeniu Spinu, Pedro Tradacete, and Vladimir G. Troitsky. Disjointly homogeneous Banach lattices: duality and complementation. *J. Funct. Anal.*, 266(9):5858–5885, 2014.
- [18] Gilles Godefroy and Nigel J. Kalton. Lipschitz-free Banach spaces. *Studia Math.*, 159(1):121–141, 2003.
- [19] Loukas Grafakos. *Classical Fourier analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014.
- [20] Richard Haydon. Injective Banach lattices. *Math. Z.*, 156(1):19–47, 1977.
- [21] Héctor Jardón-Sánchez, Niels Jakob Laustsen, Mitchell A. Taylor, Pedro Tradacete, and Vladimir G. Troitsky. Free Banach lattices under convexity conditions. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM*, 116(1):Paper No. 15, 25, 2022.
- [22] Joram Lindenstrauss and Lior Tzafriri. *Classical Banach spaces. II*, volume 97. Springer-Verlag, Berlin-New York, 1979.
- [23] Heinrich P. Lotz. Extensions and liftings of positive linear mappings on Banach lattices. *Trans. Amer. Math. Soc.*, 211:85–100, 1975.
- [24] Mieczysław Mastyło. An operator ideal generated by Orlicz spaces. *Math. Ann.*, 376(3-4):1675–1703, 2020.
- [25] Mieczysław Mastyło and Enrique A. Sánchez Pérez. Factorization of operators through Orlicz spaces. *Bull. Malays. Math. Sci. Soc.*, 40(4):1653–1675, 2017.
- [26] Mieczysław Mastyło and Radosław Szvedek. Interpolative construction and factorization of operators. *J. Math. Anal. Appl.*, 401(1):198–208, 2013.
- [27] Timur Oikhberg. Geometry of unit balls of free Banach lattices, and its applications. *J. Funct. Anal.*, to appear, 2024.
- [28] Timur Oikhberg, Mitchell A. Taylor, Pedro Tradacete, and Vladimir G. Troitsky. Free Banach lattices. *J. Eur. Math. Soc. (JEMS)*, to appear, 2024.
- [29] Timur Oikhberg and Mary Angelica Tursi. Order extreme points and solid convex hulls. In *The mathematical legacy of Victor Lomonosov – operator theory*, volume 2 of *Adv. Anal. Geom.*, pages 297–315. De Gruyter, Berlin, 2020.
- [30] Gilles Pisier. Factorization of operators through $L_{p\infty}$ or L_{p1} and noncommutative generalizations. *Math. Ann.*, 276(1):105–136, 1986.
- [31] Gilles Pisier. Complex interpolation and regular operators between Banach lattices. *Arch. Math. (Basel)*, 62(3):261–269, 1994.
- [32] Yves Raynaud and Pedro Tradacete. Interpolation of Banach lattices and factorization of p -convex and q -concave operators. *Integral Equations Operator Theory*, 66(1):79–112, 2010.
- [33] Shlomo Reisner. Operators which factor through convex Banach lattices. *Canadian J. Math.*, 32(6):1482–1500, 1980.
- [34] Shlomo Reisner. A factorization theorem in Banach lattices and its application to Lorentz spaces. *Ann. Inst. Fourier (Grenoble)*, 31(1):viii, 239–255, 1981.
- [35] Shlomo Reisner. Operators factoring through Banach lattices and ideal norms. *Serdica Math. J.*, 21(1):75–89, 1995.
- [36] Enrique A. Sánchez Pérez. Factorization theorems for multiplication operators on Banach function spaces. *Integral Equations Operator Theory*, 80(1):117–135, 2014.
- [37] Enrique A. Sánchez Pérez and Pedro Tradacete. (p, q) -regular operators between Banach lattices. *Monatsh. Math.*, 188(2):321–350, 2019.
- [38] Helmut H. Schaefer. *Banach lattices and positive operators*. Die Grundlehren der mathematischen Wissenschaften, Band 215. Springer-Verlag, New York-Heidelberg, 1974.
- [39] Vasco Schiavo. Majorizing norms on majorized spaces. *arXiv preprint arXiv:2111.06758*, 2021.
- [40] Mitchell A. Taylor and Vladimir G. Troitsky. Bibasic sequences in Banach lattices. *J. Funct. Anal.*, 278(10):108448, 33, 2020.
- [41] Lutz W. Weis. Banach lattices with the subsequence splitting property. *Proc. Amer. Math. Soc.*, 105(1):87–96, 1989.

INSTITUTO DE CIENCIAS MATEMÁTICAS (CSIC-UAM-UC3M-UCM), CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, C/ NICOLÁS CABRERA, 13–15, CAMPUS DE CANTOBLANCO UAM, 28049 MADRID, SPAIN.
Email address: `enrique.garcia@icmat.es`

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076.
Email address: `dennyhl@u.nus.edu`

DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, RAMISTRASSE 101, 8092 ZÜRICH, SWITZERLAND.
Email address: `mitchell.taylor@math.ethz.ch`

INSTITUTO DE CIENCIAS MATEMÁTICAS (CSIC-UAM-UC3M-UCM), CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, C/ NICOLÁS CABRERA, 13–15, CAMPUS DE CANTOBLANCO UAM, 28049 MADRID, SPAIN.
Email address: `pedro.tradacete@icmat.es`